

ON p -ADIC DIFFERENTIAL EQUATIONS ON SEMISTABLE VARIETIES

Valentina Di Proietto

Abstract

In this paper we prove a comparison theorem between the category of certain modules with integrable connection on the complement of a normal crossing divisor of the generic fiber of a proper semistable variety over a DVR and the category of certain log overconvergent isocrystals on the special fiber of the same open.

1 Introduction

The “théorème d’algébrisation” by Christol and Mebkhout (théorème 5.0-10 of [CtMeIV]) asserts that on a open of a smooth and proper curve all the overconvergent isocrystals (with some non-Liouville conditions) are algebraic. The aim of this paper is to give a generalized version of this result to the case of a variety of arbitrary dimension that is the special fiber of a semistable variety.

We first recall the “théorème d’algébrisation” and explain in which sense our result generalizes it. Let V be a complete discrete valuation ring of mixed characteristic $(0, p)$ with uniformizer π , let K be its fraction field and let k be the residue field. Christol and Mebkhout consider a proper and smooth curve X over V and an affine open U with complement D . They define a functor

$$\dagger : MICLS(U_K/K) \longrightarrow I^\dagger((U_k, X_k)/\mathrm{Spf}(V)) \quad (1)$$

where the first category is the subcategory of $MIC(U_K/K)$ of algebraic modules with connection on U_K which satisfy certain convergent conditions and the second is the category of overconvergent isocrystals as defined by Berthelot in [Be].

They prove that under some non-Liouville conditions the functor \dagger is essentially surjective, *i. e.* every overconvergent isocrystal is algebraic. Moreover they notice that, always assuming non-Liouville conditions, \dagger is fully faithful if one restricts to the category of algebraic modules with connection on U_K with some convergent conditions that are extendable to module with connections on X_K and logarithmic singularities along D_K . The image of this restricted functor turns out to be the category of overconvergent isocrystals with slope zero (problème 5.0-14 1) of [CtMeIV] and paragraph 6 of [CtMeII]).

We are interested in the following generalized situation.

Let X be a proper semistable variety over V , which means that locally for the étale topology is étale over $\mathrm{Spec} V[x_1, \dots, x_n, y_1, \dots, y_n]/(x_1 \cdots x_r - \pi)$ with D a normal crossing divisor, which étale locally is given by the equation $\{y_1 \cdots y_s = 0\}$. The divisor $X_k \cup D$ induces on X a logarithmic structure that we denote by M ; similarly, the closed point induces a logarithmic structure on $\mathrm{Spec}(V)$ that we denote by N . We assume we have the following diagram of fine log schemes

$$\begin{array}{ccccc} (X_k, M) & \hookrightarrow & (X, M) & \longleftarrow & (X_K, M) \\ \downarrow & & \downarrow & & \downarrow \\ (\mathrm{Spec}(k), N) & \hookrightarrow & (\mathrm{Spec}(V), N) & \longleftarrow & (\mathrm{Spec}(K), N) \end{array} \quad (2)$$

where the two squares are cartesian. The log structures denoted again by M on the special fiber X_k and by N on k are defined in such a way that the closed immersions of fine log schemes $(X_k, M) \hookrightarrow (X, M)$ and

$(\mathrm{Spec}(k), N) \hookrightarrow (\mathrm{Spec}(V), N)$ are exact. In an analogous way the log structures on X_K and on K are defined in such a way that the open immersions $(X_K, M) \hookrightarrow (X, M)$ and $(\mathrm{Spec}(K), N) \hookrightarrow (\mathrm{Spec}(V), N)$ are strict. Let us note that the log structure on K constructed in this way is isomorphic to the trivial log structure. We consider on X the open U defined as the complement of the divisor D ; then there is an open immersion

$$j : U = X \setminus D \hookrightarrow X$$

and it induces on U the log structure $j^*(M)$ that, by abuse of notation, we again denote by M . So we have a diagram analogous to (2) for U , with the same notations for the log structures:

$$\begin{array}{ccccc} (U_k, M) & \hookrightarrow & (U, M) & \longleftarrow & (U_K, M) \\ \downarrow & & \downarrow & & \downarrow \\ (\mathrm{Spec}(k), N) & \hookrightarrow & (\mathrm{Spec}(V), N) & \longleftarrow & (\mathrm{Spec}(K), N) \end{array}$$

In this situation we consider the category, denoted by $MIC(U_K/K)^{reg}$, of pairs (E, ∇) where E is a sheaf of coherent \mathcal{O}_{U_K} -modules and ∇ is an integrable connection regular along D_K .

As for the rigid side we look at $I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma}$, the category of overconvergent log isocrystals on the log pair given by $((U_k, M), (X_k, M)/(\mathrm{Spf}(V), N))$ with Σ -unipotent monodromy along D_k , where Σ is a subset of \mathbb{Z}_p^h , satisfying some non-Liouville hypothesis and h is the number of the irreducible component of D_k . A log overconvergent isocrystal is represented by a module with connection on a strict neighborhood W of the tube $]U_k[_{\hat{X}}$ in the tube $]X_k[_{\hat{X}}$. To define Σ -unipotent monodromy we proceed étale locally and we fix an irreducible component $D_{j,k}^\circ$ of D_k° , the smooth locus of D_k ; then the part of the tube $]D_{j,k}^\circ[_{\hat{X}}$ which is contained in W is isomorphic to a product of an annulus of small width times certain rigid space associate to $D_{j,k}^\circ$, that we think as a base. A log overconvergent isocrystal \mathcal{E} has Σ -unipotent monodromy along $D_{j,k}^\circ$ if \mathcal{E} , restricted to the product described above, admits a filtration such that every successive quotient is the pullback of a module with connection on the base twisted by a module with connection on the annulus depending on Σ . We say that \mathcal{E} has Σ -unipotent monodromy along D_k if the above condition holds for every irreducible component of D_k° . The category of overconvergent log isocrystals is defined by Shiho in [Sh4] as a log version of the category of overconvergent isocrystals defined by Berthelot in [Be] and the notion of Σ -unipotent monodromy is introduced by Shiho in [Sh6] as a generalization of the notion of unipotent monodromy introduced by Kedlaya in [Ke]. Let us note that the notion of Σ -unipotent monodromy for a module with connection on an annulus, with Σ satisfying some non-Liouville conditions, coincides with the notion of satisfying the Robba condition and having exponent in the sense of Christol and Mebkhout in Σ (proposition 1.18 of [Sh7]).

Our main result is:

Theorem 1. *There is a natural algebrization functor*

$$I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma} \longrightarrow MIC(U_K/K)^{reg}.$$

It is a fully faithful functor.

Let us note that our functor goes in the opposite direction with respect to Christol and Mebkhout's functor \dagger .

The strategy of our proof is as follows. We start from the category of log overconvergent isocrystals on $((U_k, M), (X_k, M)/(\mathrm{Spf}(V), N))$ with Σ -unipotent monodromy and we prove that the restriction functor

$$j^\dagger : I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf, \Sigma} \longrightarrow I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma} \quad (3)$$

is an equivalence of categories.

We denote by $I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf, \Sigma}$ the category of locally free log convergent isocrystals on the log

convergent site $((X_k, M)/(\mathrm{Spf}(V), N))_{conv}$ with exponents in Σ . The category of log convergent isocrystals is defined by Shiho in [Sh1], as a log version of the category of convergent isocrystal defined by Ogus in [Og] and by Berthelot in [Be]. The equivalence of categories in (3) is a generalization of theorem 3.16 of [Sh6], since Shiho proves the same result in the case of X_k smooth and D_k a divisor with simple normal crossing, *i. e.* its components are regular and meet transversally.

On the other hand we have a fully faithful functor

$$\tilde{i} : I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf, \Sigma} \longrightarrow MIC((X_K, M_D)/K)^{lf, \Sigma}$$

between locally free log convergent isocrystals with exponents in Σ and locally free \mathcal{O}_{X_K} -modules with integrable connection on X_K , logarithmic singularities on D_K and exponents in Σ . The theorem of algebraic logarithmic extension of [AnBa] I,4 gives us a fully faithful functor

$$MIC((X_K, M_D)/K)^{lf, \Sigma} \longrightarrow MIC(U_K/K)^{reg}. \quad (4)$$

If we denote by $MICLS(U_K/K)^{reg, \Sigma}$ the essential image of the functor \tilde{i} in $MIC(U_K/K)^{reg}$, we can conclude that we have an equivalence of categories

$$I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma} \longrightarrow MICLS(U_K/K)^{reg, \Sigma}.$$

Let us now describe in detail the contents of the paper.

In the first paragraphs we suppose that $(X, M) \rightarrow (\mathrm{Spec}(V), N)$ is a general log smooth morphism of fine log schemes (not necessarily semistable), with X proper and X_k reduced. We denote by $(\hat{X}, M) \rightarrow (\mathrm{Spf}(V), N)$ the associated morphism of the p -adic completions. We recall the definition of modules with integrable log connection on a fine log scheme, *i.e.* the category $MIC((X_K, M)/K)$, the definition of log infinitesimal isocrystals, the category $I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))$, and the definition of log convergent isocrystals, the category $I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))$. Using the fact that X is proper Shiho (Theorem 3.15, Corollary 3.2.16 of [Sh1]) shows that $MIC((X_K, M)/K)$ is equivalent to the category $I_{inf}((X_k, M)/(\mathrm{Spf}(V), N))$ of log infinitesimal isocrystals. We prove that the property of being locally free is stable with respect this equivalence of categories (propositions 17 and 19), so that we have an equivalence of categories between $I_{inf}((X_k, M)/(\mathrm{Spf}(V), N))^{lf}$ and $MIC((X_K, M)/K)^{lf}$. We use the definition of log convergent site and isocrystals on it given by Shiho in chapter 5 of [Sh1] and the functor

$$\Phi : I_{conv}(X_k, M)/(\mathrm{Spf}(V), N) \longrightarrow I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N)),$$

that he defines between log convergent isocrystals and log infinitesimal isocrystal. We adapt the proof of proposition 5.2.9 of [Sh1] to show that

$$\begin{aligned} \tilde{\Phi} : I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf} &\longrightarrow I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf} \\ &\cong MIC((X_K, M)/K)^{lf} \end{aligned} \quad (5)$$

is fully faithful (theorem 26). Shiho proves full faithfulness on the subcategory of log convergent isocrystals that are iterated extensions of the unit object. As in Shiho's proof, the key tool used is locally freeness. Finally, we give a characterization of the essential image of the functor $\tilde{\Phi}$ in terms of special stratifications, a suitable modification of the special stratifications introduced by Shiho in [Sh1]. We conclude by proving that requiring a stratification to be special is the same as requiring that the radius of convergence of the stratification is 1. In this last part we do not use log differential calculus, because we prove that we can restrict to the case of trivial log structures (proposition 28).

Then we restrict to the semistable situation: we suppose that the morphism $(X, M) \rightarrow (\mathrm{Spec}(V), N)$ is as described before (2). After describing the geometric situation, we introduce the definition of log convergent isocrystals with exponents in Σ , using log- ∇ -modules defined by Kedlaya in [Ke] and used by Shiho in [Sh6]. We first define the notion of exponents in Σ étale locally using coordinates, then we prove that it is independent of the particular étale cover chosen and from the coordinates (lemma 36). After that, following

Shiho, we recall the notion of Σ -unipotence for a $\log\text{-}\nabla$ -module defined over a product of a rigid analytic space and a polyannulus and we recall the extension theorem for Σ -unipotent $\log\text{-}\nabla$ -modules. Then we analyze log overconvergent isocrystals. Our setting is different from Shiho's ([Sh4]), since in our case the base has a non-trivial log structure. To define a log overconvergent isocrystal with Σ -unipotent monodromy we do it étale locally. Then we recall the three key propositions that we use in the proof of the main theorem. The first property is called “generalization of monodromy” and asserts that the property of being Σ -unipotent is generic on the base in some sense that we make precise (proposition 47). The second property, called “overconvergent generalization”, says that the property of being Σ -unipotent can be extended on strict neighborhoods (proposition 48). The third one says that under certain conditions a convergent $\log\text{-}\nabla$ -module that has exponents in Σ is Σ -unipotent (proposition 49). The proofs of these propositions are given in [Sh6] as a generalization the ones contained in [Ke]. Using these properties we prove that the notion of Σ -unipotent monodromy for an overconvergent isocrystal is well posed (lemma 50). Finally we prove (theorem 52) that the restriction functor

$$j^\dagger : I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf, \Sigma} \longrightarrow I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma}$$

is an equivalence of categories. The strategy of the proof is the same as in theorem 3.16 of [Sh6] and theorem 6.4.5 of [Ke].

In the last part we verify that the notion of exponents in Σ behaves well with respect to the functor \tilde{i} , *i. e.* the functor

$$I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf, \Sigma} \longrightarrow MIC((X_K, M_D)/K)^{lf, \Sigma}$$

is well defined. Finally we adapt André and Baldassarri's theorem of algebraic logarithmic extension to find theorem 1.

2 Connections with logarithmic poles

We will recall the definition of log connection on a fine log scheme or on a p -adic fine log formal scheme, which is taken from [Sh1] definition 3.1.1; see also [Kz] paragraph 4.

We suppose the reader familiar with the language of log schemes introduced in [Ka] and with his version for formal schemes given in [Sh1] chapter 2.

We denote by V a discrete valuation ring of mixed characteristic $(0, p)$, complete and separated for the p -adic topology, π a uniformizer of V , K its fraction field and k its residue field. By a formal scheme over $\mathrm{Spf}(V)$ or a formal V -scheme, we mean a p -adic Noetherian and topologically of finite type formal scheme over V . According to Shiho's notation we will give the following definition.

Definition 2. *If X is a scheme, we denote the category of coherent \mathcal{O}_X -modules by $\mathrm{Coh}(\mathcal{O}_X)$. If X is formal V -scheme we denote by $\mathrm{Coh}(K \otimes \mathcal{O}_X)$ the category of sheaves of $K \otimes_V \mathcal{O}_X$ -modules that are isomorphic to $K \otimes_V F$ for some coherent \mathcal{O}_X -module F . We will call an object of $\mathrm{Coh}(K \otimes \mathcal{O}_X)$ an isocoherent sheaf.*

The category of isocoherent sheaves is introduced in [Og] and (see [Og] remark 1.5) it is equivalent to the category of coherent sheaves on X^{an} , the rigid analytic space associated to the p -adic formal scheme X via Raynaud generic fiber.

Definition 3. *Let $f : (X, M) \rightarrow (S, N)$ be a map of fine log schemes (resp. fine log formal V -schemes) and let E be a coherent \mathcal{O}_X -module (resp. $E \in \mathrm{Coh}(K \otimes \mathcal{O}_X)$). A log connection on E is an \mathcal{O}_S -linear map*

$$\nabla : E \rightarrow E \otimes \omega_{(X, M)/(S, N)}^1$$

that is additive and satisfies the Leibniz rule:

$$\nabla(ae) = a\nabla(e) + e \otimes da$$

for $a \in \mathcal{O}_X$ and $e \in E$.

Here $\omega_{(X,M)/(S,N)}^1$ denotes the sheaf of log differentials (resp. the sheaf of log formal differentials).

We can extend ∇ to ∇_i

$$E \otimes \omega_{(X,M)/(S,N)}^i \xrightarrow{\nabla_i} E \otimes \omega_{(X,M)/(S,N)}^{i+1},$$

where $\nabla_i(e \otimes \omega) = e \otimes d\omega + (-1)^i \nabla(e) \wedge \omega$. We say that ∇ is integrable if $\nabla_{i+1} \circ \nabla_i = 0$ for all i .

We indicate the category of pairs (E, ∇) of a coherent sheaf E (resp. an isocoherent sheaf) and an integrable log connection ∇ with $MIC((X, M)/(S, N))$ (resp. $\widehat{MIC}((X, M)/(S, N))$). If M and N are isomorphic to the trivial log structures we will write $M = \text{triv}$ and $N = \text{triv}$ and we use the notation $MIC(X/S)$ (resp. $\widehat{MIC}(X/S)$) to denote $MIC((X, \text{triv})/(S, \text{triv}))$ (resp. $\widehat{MIC}((X, \text{triv})/(S, \text{triv}))$).

From proposition 8.9 of [Kz] we know that in the smooth case in characteristic zero every coherent module with integrable connection is locally free; more precisely we have the following result.

Proposition 4. *If S is the spectrum of a field of characteristic 0 and X is a smooth scheme over S , then, for every object (E, ∇) in $MIC(X/S)$, E is a locally free \mathcal{O}_X -module.*

This proposition is not true without the smoothness hypothesis; in fact, there is the following example.

Example 5. *Let X be a curve over K and D a closed point, locally defined by the equation $\{t = 0\}$. If we call M_D the log structure on X induced by D we can consider the following log connection*

$$d : \mathcal{O}_X \rightarrow \omega_{(X, M_D)/(K, \text{triv})}^1.$$

If we consider the subsheaf $\mathcal{O}_{X_K}(-D)$, that consists of sections vanishing on D , then d induces a log connection on $\mathcal{O}_{X_K}(-D)$. We can see it locally: every section of $\mathcal{O}_{X_K}(-D)$ can be written as a product of ft , with f in \mathcal{O}_{X_K} . The induced log connection is:

$$\begin{aligned} \mathcal{O}_{X_K}(-D) &\rightarrow \mathcal{O}_{X_K}(-D) \otimes \omega_{(X, M_D)/(K, \text{triv})}^1 \\ ft &\mapsto d(ft) = fdt + tdf = ftd\log t + tdf \end{aligned}$$

We can induce an integrable log connection on the the quotient $\mathcal{O}_{X_K}/\mathcal{O}_{X_K}(-D) = \mathcal{O}_D$ which is a skyscraper sheaf.

As in the case where the log structures are trivial (see for example [BeOg] chapter 1), even the category of modules with integrable log connections is equivalent to the category of log stratifications: we now describe this equivalence.

If $(X, M) \rightarrow (S, N)$ is a morphism of fine log schemes (resp. a morphism of fine log formal schemes), we denote by (X^n, M^n) the n -th log infinitesimal neighborhood of (X, M) in $(X, M) \times_{(S, N)} (X, M)$ (defined in [Ka] (5.8), [Sh1] remark 3.2.4 as a log version of the n -th infinitesimal neighborhood described in [BeOg] chapter 1).

Definition 6. *Let $(X, M) \rightarrow (S, N)$ be a morphism of fine log schemes (resp. of fine log formal V -schemes) and let E be a coherent sheaf (resp. an isocoherent sheaf on X). Then, a log stratification (resp. a formal log stratification) on E is a family of morphisms $\epsilon_n : \mathcal{O}_{X^n} \otimes E \rightarrow E \otimes \mathcal{O}_{X^n}$, that satisfy the conditions:*

- (i) ϵ_n is \mathcal{O}_{X^n} -linear and ϵ_0 is the identity;
- (ii) ϵ_n and ϵ_m are compatible via the maps

$$\mathcal{O}_{X^n} \rightarrow \mathcal{O}_{X^m}, \text{ for } m \leq n;$$

- (iii) (cocycle condition) if we call $p_{i,j}$ (for $i, j = 1, 2, 3$) the projections from the n -th log infinitesimal neighborhood $(X^n(2), M(2)^n)$ of (X, M) in $(X, M) \times_{(S, N)} (X, M) \times_{(S, N)} (X, M)$ to the n -th log infinitesimal neighborhood (X^n, M^n) of (X, M) in $(X, M) \times_{(S, N)} (X, M)$

$$p_{i,j} : (X^n(2), M^n(2)) \rightarrow (X^n, M^n),$$

then for all n

$$p_{1,2}^*(\epsilon_n) \circ p_{2,3}^*(\epsilon_n) = p_{1,3}^*(\epsilon_n).$$

We denote by $\text{Str}((X, M)/(S, M))$ the category of log stratifications (resp. with $\widehat{\text{Str}}((X, M)/(S, N))$ the category of log formal stratifications).

Theorem 3.2.15 of [Sh1] gives us the equivalence of categories that we announced: if $(X, M) \rightarrow (S, N)$ is log smooth morphism of fine log schemes over \mathbb{Q} (resp. a formally smooth morphism of fine log formal V -schemes), then

$$\begin{aligned} \text{Str}((X, M)/(S, N)) &\cong \text{MIC}((X, M)/(S, N)) \\ (\text{resp. } \widehat{\text{Str}}((X, M)/(S, N)) &\cong \widehat{\text{MIC}}((X, M)/(S, N))). \end{aligned}$$

3 Log infinitesimal isocrystals

We now define the infinitesimal site and log isocrystals on it. These are Shiho's log formal analogous of Grothendieck's infinitesimal site and crystals on it defined in [Gr] or [BeOg] chapter 1. All the definitions that follow are taken from chapter 3 of [Sh1]. We define the infinitesimal site only in the case of a morphism of fine log formal schemes, but analogous definition can be given for a morphism of log schemes.

Definition 7. Let $(\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ be a morphism of fine log formal V -schemes. An object of the log infinitesimal site $((\mathcal{X}, M)/(\mathcal{S}, N))_{\text{inf}}^{\log}$, or by brevity $(\mathcal{X}/\mathcal{S})_{\text{inf}}^{\log}$ when the log structures are clear, is a 4-ple $(\mathcal{U}, \mathcal{T}, L, \phi)$ such that \mathcal{U} is a formal V -scheme formally étale over \mathcal{X} , (\mathcal{T}, L) is a fine log formal V -scheme over (\mathcal{S}, N) and $\phi : (\mathcal{U}, M) \rightarrow (\mathcal{T}, L)$ is a nilpotent exact closed immersion of log formal V -schemes over (\mathcal{S}, N) . A morphism between $(\mathcal{U}, \mathcal{T}, L, \phi)$ and $(\mathcal{U}', \mathcal{T}', L', \phi')$ is pair of maps $g : (\mathcal{T}, L) \rightarrow (\mathcal{T}', L')$ and $f : \mathcal{U} \rightarrow \mathcal{U}'$ such that $\phi' \circ f = g \circ \phi$. The coverings in this site are the coverings of \mathcal{T} for the étale topology $\{\mathcal{T}_i \rightarrow \mathcal{T}\}_i$ such that $\mathcal{U}_i = \mathcal{T}_i \times_{\mathcal{T}} \mathcal{U}$. We sometimes denote the 4-ple $(\mathcal{U}, \mathcal{T}, L, \phi)$ simply by \mathcal{T} .

Definition 8. A log isocrystal on the infinitesimal site $(\mathcal{X}/\mathcal{S})_{\text{inf}}^{\log}$, or a log infinitesimal isocrystal, is a sheaf \mathcal{E} on $(\mathcal{X}/\mathcal{S})_{\text{inf}}^{\log}$ such that:

- (i) for every object $(\mathcal{U}, \mathcal{T}, L, \phi)$ the Zariski sheaf $\mathcal{E}_{\mathcal{T}}$ induced on \mathcal{T} is an isocoherent sheaf;
- (ii) for every morphism $g : \mathcal{T} \rightarrow \mathcal{T}'$, the map $g^*(\mathcal{E}_{\mathcal{T}'}) \rightarrow \mathcal{E}_{\mathcal{T}}$ is an isomorphism.

We denote the category of log isocrystals on the infinitesimal site $(\mathcal{X}/\mathcal{S})_{\text{inf}}^{\log}$ by $I_{\text{inf}}(\mathcal{X}/\mathcal{S})^{\log}$.

Definition 9. Let \mathcal{X} be a formal V -scheme and let \mathcal{F} be an isocoherent sheaf. We say that it is locally free module if there is a formal affine covering $\{U_i\}_{i \in I}$ of \mathcal{X} such that for every $U_i = \text{Spf} A_i$, there exists a finitely generated A_i -module F_i such that $\mathcal{F}(U_i) = K \otimes F_i$ is a projective $K \otimes A_i$ -module.

Definition 10. A log isocrystal \mathcal{E} on the infinitesimal site $(\mathcal{X}/\mathcal{S})_{\text{inf}}^{\log}$ is said to be locally free if for every object $(\mathcal{U}, \mathcal{T}, L, \phi)$ of the infinitesimal site, the sheaf $\mathcal{E}_{\mathcal{T}}$ induced on \mathcal{T} is an isocoherent locally free module. We will denote the subcategory of $I_{\text{inf}}(\mathcal{X}/\mathcal{S})^{\log}$ consisting of the locally free infinitesimal log isocrystal by $I_{\text{inf}}(\mathcal{X}/\mathcal{S})^{\log, \text{lf}}$.

Thanks to theorem 3.2.15 of [Sh1] we can see that if $(\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ is a formally log smooth morphism of fine log formal V -schemes, then there exists a canonical equivalence of categories

$$I_{\text{inf}}((\mathcal{X}, M)/(\mathcal{S}, N)) \cong \widehat{\text{MIC}}((\mathcal{X}, M)/(\mathcal{S}, N))$$

4 Log convergent isocrystals

In this section we define the log convergent site and the isocrystals on it. The following definitions are taken from [Sh1] paragraph 5.1.

Definition 11. For every log formal V -scheme (\mathcal{Y}, M) we indicate with \mathcal{Y}_1 the closed subscheme defined by the ideal p and the associated reduced subscheme of \mathcal{Y}_1 by \mathcal{Y}_0 .

Definition 12. Let $(\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ be a morphism of fine log formal V -schemes. An enlargement of (\mathcal{X}, M) over (\mathcal{S}, N) is a triple (\mathcal{T}, L, z) , that we will indicate with \mathcal{T} , such that (\mathcal{T}, L) is a fine log formal V -scheme over (\mathcal{S}, N) and z is a morphism $(\mathcal{T}_0, L) \rightarrow (\mathcal{X}, M)$ over (\mathcal{S}, N) . A morphism between two enlargements (\mathcal{T}, L, z) and (\mathcal{T}', L', z') is a morphism $g : (\mathcal{T}, L) \rightarrow (\mathcal{T}', L')$ such that $z = z' \circ g_0$, where $g_0 : (\mathcal{T}_0, L) \rightarrow (\mathcal{T}'_0, L')$ is the map induced by g .

Definition 13. We define the log convergent site of $(\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ to be the site whose objects are enlargements, morphisms are morphisms of enlargements and coverings are given by the étale topology on \mathcal{T} . We denote it by $((\mathcal{X}, M)/(\mathcal{S}, N))_{\text{conv}}$ or $(\mathcal{X}/\mathcal{S})_{\text{conv}}^{\text{log}}$ if there is no ambiguity about the log structures.

Definition 14. A log isocrystal on $(\mathcal{X}/\mathcal{S})_{\text{conv}}^{\text{log}}$, or a log convergent isocrystal, is a sheaf \mathcal{E} on $(\mathcal{X}/\mathcal{S})_{\text{conv}}^{\text{log}}$ such that:

- (i) for every enlargement (\mathcal{T}, L, z) the Zariski sheaf $\mathcal{E}_{\mathcal{T}}$ induced on \mathcal{T} is an isocoherent sheaf;
- (ii) for every morphism of enlargements $g : (\mathcal{T}, L) \rightarrow (\mathcal{T}', L')$, the map $g^*(\mathcal{E}_{\mathcal{T}'}) \rightarrow \mathcal{E}_{\mathcal{T}}$ is an isomorphism.

We denote by $I_{\text{conv}}(\mathcal{X}/\mathcal{S})^{\text{log}}$ the category of log isocrystals on the log convergent site.

Definition 15. A log isocrystal \mathcal{E} on the convergent site $((\mathcal{X}, M)/(\mathcal{S}, N))_{\text{conv}}$ is locally free if for every object \mathcal{T} on the convergent site the sheaf $\mathcal{E}_{\mathcal{T}}$ induced on \mathcal{T} is an isocoherent locally free sheaf.

We denote the subcategory of $I_{\text{conv}}(\mathcal{X}/\mathcal{S})^{\text{log}}$ consisting of the locally free log isocrystals on the convergent site by $I_{\text{conv}}(\mathcal{X}/\mathcal{S})^{\text{log, lf}}$.

5 Relations between algebraic and analytic modules with integrable connections

In what follows we consider the following situation: we fix $(X, M) \rightarrow (\text{Spec}(V), N)$ a log smooth and proper morphism of fine log schemes. We denote by $(X_K, M) \rightarrow (\text{Spec}(K), N)$ its generic fiber and $(X_k, M) \rightarrow (\text{Spec}(k), N)$ its special fiber, that we suppose to be reduced. So we have a commutative diagram

$$\begin{array}{ccccc} (X_k, M) & \hookrightarrow & (X, M) & \longleftarrow & (X_K, M) \\ \downarrow & & \downarrow f & & \downarrow \\ (\text{Spec}(k), N) & \hookrightarrow & (\text{Spec}(V), N) & \longleftarrow & (\text{Spec}(K), N) \end{array} \quad (6)$$

The log structures on X_k and $\text{Spec}(k)$, that with an abuse of notation we again call M and N , are defined in such a way that the inclusions in (X, M) and $(\text{Spec}(V), N)$ in this diagram are exact closed immersions of log schemes.

In the same way we define the log structures M on X_K and N on $\text{Spec}(K)$ as the log structures defined in such a way that the inclusions to (X, M) and $(\text{Spec}(V), N)$ are strict.

We consider the p -adic completion of $(X, M) \rightarrow (\text{Spec}(V), N)$ and we call it $(\hat{X}, M) \rightarrow (\text{Spf}(V), N)$. With (\hat{X}, M) we mean $\hat{X} = (X_k, \varprojlim \mathcal{O}_X/p^n \mathcal{O}_X)$ with the log structure, that we call again M with an abuse of notation, defined as the pull back of M via the canonical morphism $\hat{X} \rightarrow X$.

Another way to see this is [ChFo] Definition-Lemma 0.9, where the authors prove that the log structure

M over \hat{X} is isomorphic to $\varprojlim_n (M)_n$ with $(M)_n$ the log structure on $\hat{X}_n = (X_K, \mathcal{O}_X/p^n \mathcal{O}_X)$ that is the pull-back of M via the morphism $\hat{X}_n \rightarrow X$.

Now we want to construct a fully faithful functor

$$i : I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf} \longrightarrow MIC((X_K, M)/(K, N))^{lf}.$$

In corollary 3.2.16 of [Sh1] we have a useful characterization of algebraic modules with integrable log connection: the following result holds.

Proposition 16. *Under the above assumptions, there is an equivalence of categories*

$$\Psi : MIC((X_K, M)/(\mathrm{Spec}(K), N)) \longrightarrow I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N)).$$

As we saw in example 5 it is not true that every coherent module with integrable connection is locally free, so we will restrict to the category that we call

$$MIC((X_K, M)/(K, N))^{lf}$$

and that consists of pairs (E, ∇) where ∇ is an integrable log connection and E a locally free \mathcal{O}_{X_K} -module. In the next two propositions we will see that the functor Ψ of proposition 16 induces an equivalence of categories

$$MIC((X_K, M)/(K, N))^{lf} \longrightarrow I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}.$$

Proposition 17. *For every element $\mathcal{E} = (E, \nabla)$ in $MIC((X_K, M)/(K, N))^{lf}$, the corresponding element $\Psi(\mathcal{E}) \in I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))$ is a log infinitesimal locally free isocrystal.*

Proof. For the proof we look carefully at the definition of the functor Ψ . The functor is defined as the composition of three functors each one being an equivalence of category. The first functor Ψ_1 is the one that gives the equivalence of category between $MIC((X_K, M)/(K, N))$ and the category of log stratifications. So given $\mathcal{E} = (E, \nabla)$, with E a locally free \mathcal{O}_{X_K} -module, $\Psi_1(\mathcal{E})$ is again the \mathcal{O}_{X_K} -module E with a collection of isomorphisms $\epsilon_n : \mathcal{O}_{X_K^n} \otimes E \rightarrow E \otimes \mathcal{O}_{X_K^n}$, where (X_K^n, M^n) is the n -th log infinitesimal neighborhood of (X_K, M) in $(X_K, M) \times_{(K, N)} (X_K, M)$, that satisfies the conditions of definition 6. By lemma 3.2.7 of [Sh1] we know that all the $\mathcal{O}_{X_K^n}$ are free \mathcal{O}_{X_K} -modules, so $E \otimes \mathcal{O}_{X_K^n}$ are locally free \mathcal{O}_{X_K} -modules.

Now, thanks to example 1.4 of [Og], that uses a formal version of GAGA principle, we know that the category of coherent \mathcal{O}_{X_K} -modules on X_K is equivalent to the category of isocoherent modules on \hat{X} . So we can associate to our E an isocoherent sheaf \hat{E} , that we now show to be locally free. The functor that gives this equivalence is defined locally by extension of scalars: if we suppose that $X = \mathrm{Spec}(B)$ then the functor is

$$E \longmapsto E \otimes_{B_K} \hat{B}_K$$

where $B_K = B \otimes_V K$ and $\hat{B}_K = \varprojlim_n B/p^n \otimes K$. If E locally free as coherent \mathcal{O}_{X_K} -module, then E is a projective- B_K module. This implies that $E \otimes_{B_K} \hat{B}_K$ is a projective \hat{B}_K -module too. Indeed, since E is projective, there exists a B_K -module F such that $E \oplus F$ is a free B_K -module. Then $(E \oplus F) \otimes_{B_K} \hat{B}_K$ is a free \hat{B}_K -module, which implies that $E \otimes_{B_K} \hat{B}_K$ is a projective \hat{B}_K -module.

Moreover if we indicate with (\hat{X}^n, M^n) and (X_K^n, M^n) the n -th log infinitesimal neighborhoods of the diagonal morphisms $(\hat{X}, M) \rightarrow (\hat{X}, M) \times_{(\mathrm{Spf}(V), N)} (\hat{X}, M)$ and $(X_K, M) \rightarrow (X_K, M) \times_{(K, N)} (X_K, M)$ respectively, then we have an equivalence of categories also between $\mathrm{Coh}(\mathcal{O}_{X_K^n})$ and $\mathrm{Coh}(K \otimes \mathcal{O}_{\hat{X}^n})$.

This means that we have a functor Ψ_2 from the category $\mathrm{Str}((X_K, M)/K)$ to $\widehat{\mathrm{Str}}((\hat{X}, M)/(\mathrm{Spf}(V), N))$. Moreover Ψ_2 is an equivalence of categories which sends locally free objects in locally free objects.

Now we construct the functor Ψ_3 between $\widehat{\mathrm{Str}}((\hat{X}, M)/(\mathrm{Spf}(V), N))$ and $I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))$. Let us take an element $(\mathcal{U}, \mathcal{T}, L, \phi)$ of the log infinitesimal site $(\hat{X}/\mathrm{Spf}(V))_{inf}^{log}$. We know by definition that $(\mathcal{U}, M) \rightarrow (\mathrm{Spf}(V), N)$ is formally log smooth over $(\mathrm{Spf}(V), N)$, because (\mathcal{U}, M) is formally étale over

(\hat{X}, M) , that is formally log smooth over $(\mathrm{Spf} V, N)$. Therefore we have a diagram

$$\begin{array}{ccc} (\mathcal{U}, M) & \longrightarrow & (\mathcal{U}, M) \\ \phi \downarrow & & \downarrow \\ (\mathcal{T}, L) & \longrightarrow & (\mathrm{Spf}(V), N) \end{array}$$

and by proposition 2.2.13 of [Sh1] we know that étale locally over \mathcal{T} there exists a morphism $c : (\mathcal{T}, L) \rightarrow (\mathcal{U}, M)$ that is a section for $\phi : (\mathcal{U}, M) \rightarrow (\mathcal{T}, L)$.

If we have \mathcal{T}' étale over \mathcal{T} , then we call $s : (\mathcal{T}', L) \rightarrow (\hat{X}, M)$ the composition of the section c with the morphism $(\mathcal{U}, M) \rightarrow (\hat{X}, M)$.

Then we can define a sheaf over \mathcal{T}' using the pullback map $s_K^* : \mathrm{Coh}(K \otimes \mathcal{O}_{\hat{X}}) \rightarrow \mathrm{Coh}(K \otimes \mathcal{O}_{\mathcal{T}'})$, and we call the resulting sheaf $\hat{E}_{\mathcal{T}'} = s_K^* \hat{E}$. Let us note that $E_{\mathcal{T}'}$ is obviously a locally free isocoherent sheaf on \mathcal{T}' , as soon as \hat{E} is.

If we have two sections c and d and respectively two morphisms s and t , the formally log smooth morphism $s \times t : (\mathcal{T}', L) \rightarrow (\hat{X}, M) \times_{(\mathrm{Spf}(V), N)} (\hat{X}, M)$ factors through the n -th log infinitesimal neighborhood (\hat{X}^n, M^n) , for some n :

$$s \times t : (\mathcal{T}', L) \xrightarrow{u} (\hat{X}^n, M^n) \longrightarrow (\hat{X}, M) \times_{(\mathrm{Spf}(V), N)} (\hat{X}, M),$$

because of the universal propriety of the n -th infinitesimal neighborhood.

Now, pulling back by u the isomorphisms $\hat{e}_n : \mathcal{O}_{\hat{X}^n} \otimes \hat{E} \rightarrow \hat{E} \otimes \mathcal{O}_{\hat{X}^n}$ given by the stratification, we obtain an isomorphism $t_K^* \hat{E} \cong s_K^* \hat{E}$. We want to descend this sheaf to \mathcal{T} ; this is possible using the cocycle condition and the theorem of faithfully flat descent for isocoherent sheaves of Gabber ([Og] proposition 1.9.).

In fact, let us consider the formally étale morphism $(\mathcal{T}', L) \rightarrow (\mathcal{T}, L)$ and the two projections

$$\begin{array}{ccc} (\mathcal{T}', L) \times_{(\mathcal{T}, L)} (\mathcal{T}', L) & \xrightarrow{p_2} & (\mathcal{T}', L) \\ p_1 \downarrow & & \\ (\mathcal{T}', L) & & \end{array} .$$

Composing p_i with the morphism $s : (\mathcal{T}', L) \rightarrow (\hat{X}, M)$ we find $\pi_i : (\mathcal{T}', L) \times_{(\mathcal{T}, L)} (\mathcal{T}', L) \rightarrow (\hat{X}, M)$ for $i = 1, 2$ respectively.

As before we have a map $\pi_1 \times \pi_2 : (\mathcal{T}', L) \times_{(\mathcal{T}, L)} (\mathcal{T}', L) \rightarrow (\hat{X}, M) \times_{(\mathrm{Spf}(V), N)} (\hat{X}, M)$ that factors through the n -th log infinitesimal neighborhood and we can deduce an isomorphism $\pi_{1,K}^* \hat{E} \cong \pi_{2,K}^* \hat{E}$ and consequently an isomorphism $p_{1,K}^* \hat{E}_{\mathcal{T}'} \cong p_{2,K}^* \hat{E}_{\mathcal{T}'}$, that is a covering datum; to obtain a descent datum we adapt the above argument using the cocycle condition of the stratification. Now using proposition 1.9 of [Og] we are allowed to descend the sheaf $\hat{E}_{\mathcal{T}'}$ to an isocoherent sheaf on \mathcal{T} , that we call $\hat{E}_{\mathcal{T}}$.

Let us note that $\hat{E}_{\mathcal{T}}$ defines a log isocrystal on the infinitesimal site: to check property (ii) of definition 8 we can use the same arguments as before.

Now we prove that $\hat{E}_{\mathcal{T}}$ is a locally free isocoherent module on \mathcal{T} .

We know that $\hat{E}_{\mathcal{T}}$ is an isocoherent module on \mathcal{T} , isomorphic to $K \otimes F$, for some coherent sheaf F of $\mathcal{O}_{\mathcal{T}}$ -modules, and that there exists an étale covering of \mathcal{T} such that the $\hat{E}_{\mathcal{T}}$, restricted to every element of the covering, is a locally free isocoherent module. We can restrict ourselves to the affine case; we assume that $\mathcal{T} = \mathrm{Spf}(A)$, $\mathcal{T}' = \mathrm{Spf}(B)$ and $\mathrm{Spf}(B) \rightarrow \mathrm{Spf}(A)$ étale surjective. We can conclude applying the following lemma. \square

Lemma 18. *Let A and B be commutative noetherian V -algebras and let M be a finitely generated $A \otimes K = A_K$ -module. If we have a map $A \rightarrow B$ that is faithfully flat and $B_K \otimes_{A_K} M$ is projective, then M is a projective A_K -module.*

Proof. We have the following isomorphism for every A_K -module N :

$$B_K \otimes_{A_K} \mathrm{Ext}_{A_K}^i(M, N) \cong \mathrm{Ext}_{B_K}^i(B_K \otimes_{A_K} M, B_K \otimes_{A_K} N),$$

for every $i \geq 0$, because B_K is flat over A_K . We can conclude that M is projective because $B_K \otimes_{A_K} M$ is assumed to be projective. \square

It is true also the viceversa of proposition 17.

Proposition 19. *If \mathcal{E} is a log infinitesimal locally free isocrystal, then there exists an object $(E, \nabla) \in MIC((X_K, M)/(K, N))^{lf}$ such that $\Psi((E, \nabla)) = \mathcal{E}$.*

Proof. From proposition 16 we know that there exists an element (E, ∇) in $MIC((X_K, M)/(K, N))$ such that $\Psi((E, \nabla)) = \mathcal{E}$.

So we have to show that E is a locally free \mathcal{O}_{X_K} -module. By proposition 17 we are reduced to prove that the equivalence of categories

$$j : \text{Coh}(\mathcal{O}_{X_K}) \rightarrow \text{Coh}(K \otimes \mathcal{O}_{\hat{X}}) \quad (7)$$

behaves well with respect to locally free objects, in particular that if \mathcal{F} is a locally free isocoherent module then there exists a locally free sheaf of \mathcal{O}_{X_K} -modules F such that $j(F) = \mathcal{F}$.

Let us take $F \in \text{Coh}(\mathcal{O}_{X_K})$ such that $\mathcal{F} = j(F)$. We claim that if F is not locally free, then \mathcal{F} is not locally free. Let us assume that F is not locally free. So there exists an open affine, that we can suppose local, $U = \text{Spec}(A) \subseteq X_K$ such that $F|_U$ is not flat on U . By definition of flatness there exists a coherent ideal \mathcal{I} of A such that $\mathcal{I} \otimes F|_U \rightarrow F|_U$ is not injective. Let us take a coherent ideal \mathcal{I}' of \mathcal{O}_{X_K} that extends \mathcal{I} ([Ha] ex. II 5.15). Then $\mathcal{I}' \otimes F \rightarrow F$ is not injective. Therefore, since the functor j is faithful, exact and compatible with tensor products the map $j(\mathcal{I}') \otimes \mathcal{F} = j(\mathcal{I}' \otimes F) \rightarrow j(F) = \mathcal{F}$ is not injective. So the functor

$$- \otimes \mathcal{F} : \text{Coh}(K \otimes \mathcal{O}_{\hat{X}}) \rightarrow \text{Coh}(K \otimes \mathcal{O}_{\hat{X}})$$

is not an exact functor and then \mathcal{F} is not locally free. \square

Now, as in [Sh1] paragraph 5.2, we construct a functor

$$\Phi : I_{conv}((\hat{X}, M)/(\text{Spf}(V), N)) \longrightarrow I_{inf}((\hat{X}, M)/(\text{Spf}(V), N)).$$

We remark that the functor Φ and its restriction to locally free objects, $\tilde{\Phi}$, that we will mention below, can be constructed for $(\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$, morphism of fine log formal schemes.

Let $(\mathcal{U}, \mathcal{T}, L, \phi)$ be an object of the infinitesimal site, define an enlargement $\Phi^*(\mathcal{T}) = (\mathcal{T}, L, z : (\mathcal{T}_0, L) \cong (\mathcal{U}_0, M) \rightarrow (\hat{X}, M))$; this is clearly an element of the log convergent site. Let us observe that $\Phi^*(\hat{X}) = (\hat{X}, M, z : (X_k, M) \rightarrow (\hat{X}, M))$.

If we have an isocrystal \mathcal{E} on the log convergent site we define

$$\Phi(\mathcal{E})_{\mathcal{T}} = \mathcal{E}_{\Phi^*(\mathcal{T})}.$$

We have already saw in proposition 17 and 19 that $\tilde{\Psi}$, the restriction of the functor Ψ to the locally free objects

$$\tilde{\Psi} : MIC((X_K, M)/(\text{Spf}(V), N))^{lf} \longrightarrow I_{inf}((\hat{X}, M)/(\text{Spf}(V), N))^{lf},$$

is well-defined and is an equivalence of categories. Now we want to show that also $\tilde{\Phi}$, the restriction of the functor Φ to the locally free objects

$$\tilde{\Phi} : I_{conv}((\hat{X}, M)/(\text{Spf}(V), N))^{lf} \longrightarrow I_{inf}((\hat{X}, M)/(\text{Spf}(V), N))^{lf},$$

is well-defined. In particular we have the following lemma.

Proposition 20. *If \mathcal{E} is a log convergent isocrystal, such that $\Phi(\mathcal{E})$ is a locally free log infinitesimal isocrystal, then also \mathcal{E} is locally free.*

Proof. We can evaluate \mathcal{E} at the enlargement $(\hat{X}, M, z : (X_k, M) \rightarrow (\hat{X}, M))$ and we have that $\mathcal{E}_{\hat{X}} = \mathcal{E}_{\Phi^*(\hat{X})} = \Phi(E)_{\hat{X}}$; so $\mathcal{E}_{\hat{X}}$ is a locally free isocoherent module. Now taken an enlargement $(\mathcal{T}, L, z : (\mathcal{T}_0, L) \rightarrow (\hat{X}, M))$, we have the following commutative diagram of log formal schemes

$$\begin{array}{ccc} (\mathcal{T}_0, L) & \xrightarrow{z} & (\hat{X}, M) \\ \downarrow i & & \downarrow f \\ (\mathcal{T}, L) & \xrightarrow{b} & (\mathrm{Spf}(V), N). \end{array}$$

As f is formally log smooth, we know that étale locally on \mathcal{T} there is a morphism $c : (\mathcal{T}, L) \rightarrow (\hat{X}, M)$ such that $c \circ i = z$ and $f \circ c = b$.

So let us consider \mathcal{T}' formally étale over \mathcal{T} and the enlargement $\mathcal{T}' = (\mathcal{T}', L, z : (\mathcal{T}'_0, L) \rightarrow (\hat{X}, M))$, and call again c the morphism $(\mathcal{T}', L) \rightarrow (\hat{X}, M)$; this clearly extends to a morphism of enlargements. So by definition of isocrystal we have an isomorphism $c^*(\mathcal{E}_{\hat{X}}) \cong \mathcal{E}_{\mathcal{T}'}$, from which we know that $\mathcal{E}_{\mathcal{T}'}$ is a locally free isocoherent module.

Then we consider the formally étale morphism $u : (\mathcal{T}', L) \rightarrow (\mathcal{T}, L)$ that extends to a morphism of enlargements; again by definition of isocrystal we have $u^*(\mathcal{E}_{\mathcal{T}}) \cong \mathcal{E}_{\mathcal{T}'}$. We know that $\mathcal{E}_{\mathcal{T}'}$ is a locally free isocoherent module and we want to prove that $\mathcal{E}_{\mathcal{T}}$ is locally free: we can proceed as in the last part of proposition 17 and conclude. \square

From the definition of the functor Φ we see also the viceversa of proposition 20; indeed if \mathcal{E} is a locally free log convergent isocrystal, then for every element \mathcal{T} on the log infinitesimal site

$$\Phi(\mathcal{E})_{\mathcal{T}} = \mathcal{E}_{\Phi^*(\mathcal{T})},$$

by definition of Φ .

So we can compose the functors $\tilde{\Phi}^{-1}$ and $\tilde{\Psi}^{-1}$ and obtain a well defined functor

$$i : I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf} \rightarrow MIC((X_K, M)/(\mathrm{Spf}(V), N))^{lf}.$$

Our next goal is to prove that $\tilde{\Phi}$ is fully faithful, showing first that this can be proved étale locally.

Proposition 21. *If $\tilde{\Phi} : I_{conv}(\hat{X}/\mathrm{Spf}(V))^{log,lf} \rightarrow I_{inf}(\hat{X}/\mathrm{Spf}(V))^{log,lf}$ is fully faithful étale locally on \hat{X} , then $\tilde{\Phi} : I_{conv}(\hat{X}/\mathrm{Spf}(V))^{log,lf} \rightarrow I_{inf}(\hat{X}/\mathrm{Spf}(V))^{log,lf}$ is fully faithful.*

Proof. We suppose that $\coprod_j \hat{X}_j = \hat{X}^e \rightarrow \hat{X}$ is an étale covering of \hat{X} . If $\hat{X}' = \hat{X}^e \times_{\hat{X}} \hat{X}^e$ and $\hat{X}'' = \hat{X}^e \times_{\hat{X}} \hat{X}^e \times_{\hat{X}} \hat{X}^e$, we have the following diagram:

$$\begin{array}{ccc} I_{conv}(\hat{X}/\mathrm{Spf}(V))^{log,lf} & \xrightarrow{\tilde{\Phi}} & I_{inf}(\hat{X}/\mathrm{Spf}(V))^{log,lf} \\ \downarrow & & \downarrow \\ I_{conv}(\hat{X}^e/\mathrm{Spf}(V))^{log,lf} & \xrightarrow{\tilde{\Phi}^e} & I_{inf}(\hat{X}^e/\mathrm{Spf}(V))^{log,lf} \\ \Downarrow & & \Downarrow \\ I_{conv}(\hat{X}'/\mathrm{Spf}(V))^{log,lf} & \xrightarrow{\tilde{\Phi}'} & I_{inf}(\hat{X}'/\mathrm{Spf}(V))^{log,lf} \\ \Downarrow & & \Downarrow \\ I_{conv}(\hat{X}''/\mathrm{Spf}(V))^{log,lf} & \xrightarrow{\tilde{\Phi}''} & I_{inf}(\hat{X}''/\mathrm{Spf}(V))^{log,lf} \end{array}$$

where the vertical arrows are induced by the following morphisms:

$$\hat{X}^e \rightarrow \hat{X},$$

$$p_i : \hat{X}' = \hat{X}^e \times_{\hat{X}} \hat{X}^e \rightarrow \hat{X}^e \text{ for } i = 1, 2,$$

$$p_{i,j} : \hat{X}'' = \hat{X}^e \times_{\hat{X}} \hat{X}^e \times_{\hat{X}} \hat{X}^e \rightarrow \hat{X}^e \times_{\hat{X}} \hat{X}^e \text{ for } 1 \leq i < j \leq 3$$

where the first is the étale morphism defining the étale cover, p_i and $p_{i,j}$ are the natural projections.

Thanks to étale descent for log convergent isocrystals ([Sh1] remark 5.1.7), giving \mathcal{E} , a locally free log convergent isocrystal on \hat{X} is the same as giving \mathcal{E}^e , a locally free log convergent isocrystal on \hat{X}^e and $\alpha_{\mathcal{E}}$ isomorphism between $p_2^* \mathcal{E}^e \rightarrow p_1^* \mathcal{E}^e$, compatible with the usual cocycle conditions.

A morphism f from \mathcal{E} to \mathcal{F} is the same as a morphism f^e from \mathcal{E}^e and \mathcal{F}^e that satisfies $\alpha_{\mathcal{F}} \circ p_2^* f^e = p_1^* f^e \circ \alpha_{\mathcal{E}}$ and the compatibility conditions given by cocycle conditions on \hat{X}'' .

By hypothesis $\tilde{\Phi}^e$, $\tilde{\Phi}'$ and $\tilde{\Phi}''$ are fully faithful, so f^e induces a unique morphism $\tilde{\Phi}^e(f^e)$ between $\tilde{\Phi}^e(\mathcal{E}^e) \rightarrow \tilde{\Phi}^e(\mathcal{F}^e)$ that satisfies the same compatibility conditions on \hat{X}' and \hat{X}'' . Moreover this association is surjective. Using étale descent for log infinitesimal isocrystals proven in [Sh1] remark 3.2.20 we can descend $\tilde{\Phi}^e(f^e)$ to a morphism which coincides with $\tilde{\Phi}(f)$ between $\tilde{\Phi}(\mathcal{E})$ and $\tilde{\Phi}(\mathcal{F})$. \square

Before proving the full faithfulness étale locally we recall the construction of the universal enlargement and of convergent stratifications given by Shiho in [Sh1] paragraph 5.2.

Let $\ell : (\mathcal{X}, M) \hookrightarrow (\mathcal{Y}, M')$ be an exact closed immersion of p -adic log formal V -schemes over (\mathcal{S}, N) defined by a sheaf of ideals \mathcal{I} . We need a more general notion of enlargement.

Definition 22. We say that the quadruple (\mathcal{T}, L, z, g) is an enlargement of (\mathcal{X}, M) in (\mathcal{Y}, M') if (\mathcal{T}, L, z) is an enlargement of (\mathcal{X}, M) and g is an (\mathcal{S}, N) morphism $(\mathcal{T}, L) \rightarrow (\mathcal{Y}, M')$ such that the following diagram is commutative

$$\begin{array}{ccc} (\mathcal{T}_0, L) & \xrightarrow{g_0} & (\mathcal{Y}_0, M') \\ \downarrow z & & \downarrow \\ (\mathcal{X}, M) & \xrightarrow{\ell} & (\mathcal{Y}, M') \end{array}$$

We call $\mathcal{B}_{n,\mathcal{X}}(\mathcal{Y})$ the formal blow up of (\mathcal{Y}, M') with respect to $p\mathcal{O}_{\mathcal{Y}} + \mathcal{I}^{n+1}$ and we denote by $T_{n,\mathcal{X}}(\mathcal{Y})$ the open of $\mathcal{B}_{n,\mathcal{X}}(\mathcal{Y})$ defined by p :

$$T_{n,\mathcal{X}}(\mathcal{Y}) := \{x \in \mathcal{B}_{n,\mathcal{X}}(\mathcal{Y}) \mid (p\mathcal{O}_{\mathcal{Y}} + \mathcal{I}^{n+1})\mathcal{O}_{\mathcal{B}_{n,\mathcal{X}}(\mathcal{Y}),x} = p\mathcal{O}_{\mathcal{B}_{n,\mathcal{X}}(\mathcal{Y}),x}\}.$$

We put on $T_{n,\mathcal{X}}(\mathcal{Y})$ the log structure $L_{n,\mathcal{X}}(\mathcal{Y})$ induced by the pull-back of M' .

Remark 23. We can see from [Og] proposition 2.3 that if $\mathcal{Y} = \mathrm{Spf}(A)$ and $\mathcal{I}^{n+1} = (a_1, \dots, a_m)$ then $T_{n,\mathcal{X}}(\mathcal{Y}) = \mathrm{Spf}(A\{t_1, \dots, t_m\}/(pt_1 - a_1, \dots, pt_m - a_m))$ modulo p -torsion. For a local description of the formal blow up see [Bo] paragraph 2.6.

The map $(T_{n,\mathcal{X}}(\mathcal{Y}))_0 \rightarrow (\mathcal{Y})_0$ factors through \mathcal{X}_0 , so that we can equip the pair $(T_{n,\mathcal{X}}(\mathcal{Y}), L_{n,\mathcal{X}}(\mathcal{Y}))$ with two maps (z_n, t_n) in such a way that the quadruple $((T_{n,\mathcal{X}}(\mathcal{Y}), L_{n,\mathcal{X}}(\mathcal{Y}), z_n, t_n)$ is an enlargement of (\mathcal{X}, M) in (\mathcal{Y}, M') and the set $\{(T_{n,\mathcal{X}}(\mathcal{Y}), L_{n,\mathcal{X}}(\mathcal{Y}), z_n, t_n)\}_{n \in \mathbb{N}}$ is an inductive system of enlargements that is universal in the sense that for every enlargement (T', L', z', t') of (\mathcal{X}, M) in (\mathcal{Y}, M') there exists a unique morphism to the inductive system given by $\{(T_{n,\mathcal{X}}(\mathcal{Y}), L_{n,\mathcal{X}}(\mathcal{Y}), z_n, t_n)\}_{n \in \mathbb{N}}$ (proposition 5.2.4 of [Sh1]).

The fiber product of fine log formal schemes is a log formal scheme that is not necessarily fine, but, thanks to proposition 2.1.6 of [Sh1], there is a functor $(-)^{int}$ that sends a log formal scheme to a fine log formal scheme that is right adjoint to the natural inclusion of fine log formal schemes in the category of log formal schemes. If $f : (\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ is a morphism of fine log formal schemes, we denote the fiber product in the category of log formal schemes of (\mathcal{X}, M) and (\mathcal{X}, M) over (\mathcal{S}, N) by $(\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M)$ (resp. the fiber product in the category of log formal schemes of (\mathcal{X}, M) with itself three times over (\mathcal{S}, N) with $(\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M)$) and the fine log formal scheme associated to this with $((\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M))^{int}$ (resp. $((\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M))^{int}$).

We want to construct a log formal scheme that we will indicated by $(\mathcal{X}(1), M(1))$ (resp. $(\mathcal{X}(2), M(2))$) which

factors the diagonal embedding $\Delta^{int} : (\mathcal{X}, M) \rightarrow ((\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M))^{int}$ (resp. $\Delta^{int} : (\mathcal{X}, M) \rightarrow ((\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M))^{int}$) in an exact closed immersion $(\mathcal{X}, M) \hookrightarrow (\mathcal{X}(1), M(1))$ (resp. the closed immersion $(\mathcal{X}, M) \hookrightarrow (\mathcal{X}(2), M(2))$) followed by a formally log étale morphism: following [Sh1] or [Ka] proposition 4.10, we can do this if the morphism $f : (\mathcal{X}, M) \rightarrow (\mathcal{S}, N)$ has a chart.

We indicate with $(P_{\mathcal{X}} \rightarrow M, Q_V \rightarrow N, Q \rightarrow P)$ a chart of f , with $\alpha(1)$ (resp. $\alpha(2)$) the homomorphism induced by the map $P \oplus_Q P \rightarrow P$ (resp. $P \oplus_Q P \oplus_Q P \rightarrow P$) and with $R(1)$ the set $(\alpha(1)^{gp})^{-1}(P)$ (resp. with $R(2)$ the set $(\alpha(2)^{gp})^{-1}(P)$).

With this notation we define $\mathcal{X}(1) = (\mathcal{X} \times_{\mathcal{S}} \mathcal{X}) \times_{\mathrm{Spf}(\mathbb{Z}_p\{P \oplus_Q P\})} \mathrm{Spf}(\mathbb{Z}_p\{R(1)\})$ (resp. $\mathcal{X}(2) = (\mathcal{X} \times_{\mathcal{S}} \mathcal{X} \times_{\mathcal{S}} \mathcal{X}) \times_{\mathrm{Spf}(\mathbb{Z}_p\{P \oplus_Q P \oplus_Q P\})} \mathrm{Spf}(\mathbb{Z}_p\{R(2)\})$) equipped with the log structure $M(1)$ (resp $M(2)$) defined as the log structure induced by the canonical log structure on $\mathrm{Spf}(\mathbb{Z}_p\{R(1)\})$ (resp. on $\mathrm{Spf}(\mathbb{Z}_p\{R(2)\})$). Thanks to proposition 4.10 of [Ka] we have that $(\mathcal{X}(1), M(1))$ (resp. $(\mathcal{X}(2), M(2))$) factors the diagonal embedding as we wanted.

Using the fact that $(\mathcal{X}, M) \hookrightarrow (\mathcal{X}(i), M(i))$ are exact closed immersions of log formal schemes for $i = 1, 2$, we define $\{(T_{\mathcal{X},n}(\mathcal{X}(i)), L_{\mathcal{X},n}(\mathcal{X}(i)), z_n(i), t_n(i))\}_{n \in \mathbb{N}}$ which is the universal system of enlargements associated to this closed immersions. For simplicity of notation we will denote by $(T_n(i), L_n(i))$ the n -th universal enlargement $(T_{\mathcal{X},n}(\mathcal{X}(i)), L_{\mathcal{X},n}(\mathcal{X}(i)), z_n(i), t_n(i))$.

The natural maps

$$\begin{aligned} p_i &: (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M) \rightarrow (\mathcal{X}, M) & \text{for } i = 1, 2, \\ p_{i,j} &: (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, N) \times_{(\mathcal{S}, N)} (\mathcal{X}, N) \rightarrow (\mathcal{X}, N) \times_{(\mathcal{S}, N)} (\mathcal{X}, N) \\ & \text{for } 1 \leq i < j \leq 3 \end{aligned}$$

$$\Delta : (\mathcal{X}, M) \rightarrow (\mathcal{X}, M) \times_{(\mathcal{S}, N)} (\mathcal{X}, M)$$

induce compatible morphisms of enlargements:

$$\begin{aligned} q_{i;n} &: (T_n(1), L_n(1)) \rightarrow (\mathcal{X}, M) \\ q_{i,j;n} &: (T_n(2), L_n(1)) \rightarrow (T_n(1), L_n(1)) \\ \Delta_n &: (\mathcal{X}, M) \rightarrow (T_n(1), L_n(1)). \end{aligned}$$

With the same notation we can give the following definition.

Definition 24. A log convergent stratification on (\mathcal{X}, M) is an isocoherent sheaf $\mathcal{E}_{\mathcal{X}}$ on \mathcal{X} and a compatible family of isomorphisms

$$\epsilon_n : q_{2;n}^* \mathcal{E}_{\mathcal{X}} \rightarrow q_{1;n}^* \mathcal{E}_{\mathcal{X}}$$

such that every ϵ_n satisfies

$$\begin{aligned} \Delta_n^*(\epsilon_n) &= id; \\ q_{1,2;n}^*(\epsilon_n) \circ q_{2,3;n}^*(\epsilon_n) &= q_{1,3;n}^*(\epsilon_n). \end{aligned}$$

We denote the category of log convergent stratifications by $\mathrm{Str}'((\mathcal{X}, M)/(\mathcal{S}, N))$.

As in the case of log infinitesimal isocrystals we can establish an equivalence of categories between log convergent stratifications and convergent log isocrystals: this is the statement of proposition 5.2.6 of [Sh1].

Proposition 25. If the log formal scheme (\mathcal{X}, M) is formally log smooth over the log formal scheme $(\mathrm{Spf}(V), N)$, then the category $I_{conv}((\mathcal{X}, M)/(\mathrm{Spf}(V), N))$ is equivalent to $\mathrm{Str}'((\mathcal{X}, M)/(\mathrm{Spf}(V), N))$.

Now that we have introduced all the machinery we can finish the proof of full faithfulness of the functor $\tilde{\Phi}$. We need the following result, whose proof is essentially the same as the one of proposition 5.2.9 of [Sh1].

Theorem 26. The functor

$$\tilde{\Phi} : I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf} \longrightarrow I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$$

is fully faithful étale locally.

Proof. Since fine log formal schemes have charts étale locally and the statement is of étale local nature, we can suppose that the morphism $f : (\hat{X}, M) \rightarrow (\mathrm{Spf}(V), N)$ has a chart globally. Let us note first that for \mathcal{E} and \mathcal{F} in $I_{\mathrm{conv}}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$, we can define $\mathcal{H}om(\mathcal{E}, \mathcal{F}) \in I_{\mathrm{conv}}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$ by

$$\mathcal{H}om(\mathcal{E}, \mathcal{F})_{\mathcal{T}} = \mathcal{H}om(\mathcal{E}_{\mathcal{T}}, \mathcal{F}_{\mathcal{T}})$$

and moreover that the global sections of $\mathcal{H}om(\mathcal{E}, \mathcal{F})$ are isomorphic to $\mathrm{Hom}(\mathcal{E}, \mathcal{F})$; these two assertions follow from the fact that \mathcal{E} and \mathcal{F} are locally free.

Using the local freeness we can conclude that the same holds for the category $I_{\mathrm{inf}}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$. So we are reduced to prove that there is an isomorphism

$$H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{conv}}, \mathcal{E}) \longrightarrow H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{inf}}, \tilde{\Phi}(\mathcal{E}))$$

for every \mathcal{E} in $I_{\mathrm{conv}}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$.

As we noticed before the morphism $f : (\hat{X}, M) \rightarrow (\mathrm{Spf}(V), N)$ has a chart globally and so we can construct the scheme $(\hat{X}(1), (M)(1))$ that we described above.

The equivalence in proposition 25 associates to $\mathcal{E} \in I_{\mathrm{conv}}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$ a log convergent stratification $(\mathcal{E}_{\hat{X}}, \epsilon_n)$ given by a locally free isocoherent sheaf $\mathcal{E}_{\hat{X}}$ on \hat{X} and isomorphisms

$$\epsilon_n : (K \otimes \mathcal{O}_{T_n(1)}) \otimes \mathcal{E}_{\hat{X}} \longrightarrow \mathcal{E}_{\hat{X}} \otimes (K \otimes \mathcal{O}_{T_n(1)})$$

for all n .

So the set $H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{conv}}, \mathcal{E})$ can be characterized in terms of log convergent stratifications as follows:

$$H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{conv}}, \mathcal{E}) = \{ e \in \Gamma(\hat{X}, \mathcal{E}_{\hat{X}}) \mid \epsilon_n(1 \otimes e) = e \otimes 1 \ \forall n \}.$$

Let J be the sheaf of ideals that defines the closed immersion $\hat{X} \hookrightarrow \hat{X}(1)$; we denote by $\mathcal{O}_{\hat{X}(1)^{\mathrm{an}}}$ the sheaf $\varprojlim_m K \otimes \mathcal{O}_{\hat{X}(1)}/J^m$. By proposition 5.2.7 (2) of [Sh1] we know that there is an injective map $K \otimes \mathcal{O}_{T_n(1)} \rightarrow \mathcal{O}_{\hat{X}(1)^{\mathrm{an}}}$. If we tensor the isomorphisms ϵ_n of the convergent stratification $(\mathcal{E}_{\hat{X}}, \epsilon_n)$ with this map we obtain a map

$$\epsilon' : \mathcal{O}_{\hat{X}(1)^{\mathrm{an}}} \otimes \mathcal{E}_{\hat{X}} \longrightarrow \mathcal{E}_{\hat{X}} \otimes \mathcal{O}_{\hat{X}(1)^{\mathrm{an}}}$$

that coincides with the limit of the isomorphisms of the stratification induced by $\tilde{\Phi}(\mathcal{E})$ through the equivalence of categories

$$I_{\mathrm{inf}}((\hat{X}, M)/(\mathrm{Spf}(V), N)) \cong \widehat{\mathrm{Str}}((\hat{X}, M)/(\mathrm{Spf}(V), N)).$$

So we can characterize the set $H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{inf}}, \tilde{\Phi}(\mathcal{E}))$ as follows

$$H^0((\hat{X}, M)/(\mathrm{Spf}(V), N)_{\mathrm{inf}}, \tilde{\Phi}(\mathcal{E})) = \{ e \in \Gamma(\hat{X}, \mathcal{E}_{\hat{X}}) \mid \epsilon'(1 \otimes e) = e \otimes 1 \}.$$

This means that the claim is reduced to prove that the following diagram

$$\begin{array}{ccc} \mathcal{O}_{\hat{X}(1)^{\mathrm{an}}} \otimes \mathcal{E}_{\hat{X}} & \xrightarrow{\epsilon'} & \mathcal{E}_{\hat{X}} \otimes \mathcal{O}_{\hat{X}(1)^{\mathrm{an}}} \\ & \nwarrow \quad \nearrow & \\ & \mathcal{E}_{\hat{X}} & \end{array} \quad (8)$$

is commutative if and only if this is commutative

$$\begin{array}{ccc} (K \otimes \mathcal{O}_{T_n(1)}) \otimes \mathcal{E}_{\hat{X}} & \xrightarrow{\epsilon_n} & \mathcal{E}_{\hat{X}} \otimes (K \otimes \mathcal{O}_{T_n(1)}) \\ & \nwarrow \quad \nearrow & \\ & \mathcal{E}_{\hat{X}} & \end{array} \quad (9)$$

Knowing that the following diagram is commutative

$$\begin{array}{ccc}
\mathcal{O}_{\hat{X}(1)^{\text{an}}} \otimes \mathcal{E}_{\hat{X}} & \xrightarrow{\epsilon'} & \mathcal{E}_{\hat{X}} \otimes \mathcal{O}_{\hat{X}(1)^{\text{an}}} \\
\uparrow & & \uparrow \\
(K \otimes \mathcal{O}_{T_n(1)}) \otimes \mathcal{E}_{\hat{X}} & \xrightarrow{\epsilon_n} & \mathcal{E}_{\hat{X}} \otimes (K \otimes \mathcal{O}_{T_n(1)})
\end{array} \tag{10}$$

and putting together (9) and (10), we can conclude that if (9) is commutative then (8) is commutative. Let us suppose instead that (8) is commutative: then using the fact that $\mathcal{E}_{\hat{X}}$ is flat and that the map $K \otimes \mathcal{O}_{T_n(1)} \rightarrow \mathcal{O}_{\hat{X}(1)^{\text{an}}}$ is injective we can conclude that also (9) is commutative. \square

Shiho in [Sh1] proposition 5.2.9 proves that the functor Φ is fully faithful étale locally when it is restricted to the nilpotent part of $I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))$ and $I_{\text{inf}}((\hat{X}, M)/(\text{Spf}(V), N))$. Our proof is essentially the same, because the key property of nilpotent objects used in Shiho's proof is that the nilpotent isocrystals are locally free.

Putting together theorem 26 and proposition 21 we obtain the following

Theorem 27. *The functor $\tilde{\Phi}$ is fully faithful.*

6 Characterizations of log convergent isocrystals in terms of stratifications

We want to describe the essential image of the functor $\tilde{\Phi}$. As for the case of the proof of full faithfulness it will be enough to describe it étale locally, then, with descent argument we can conclude as in proposition 21. So we can suppose that $(\hat{X}, M) \rightarrow (\text{Spf}(V), N)$ has a chart globally.

To avoid log differential calculus we will prove that we can restrict to the case of trivial log structures. We need the following lemma whose proof is essentially the same as proposition 5.2.11 of [Sh1].

Proposition 28. *Let $f : (\hat{X}, M) \rightarrow (\text{Spf}(V), N)$ be a formally smooth morphism of fine log schemes that admits a chart. Let U be a dense open subset of \hat{X} and set j the open immersion $j : U \hookrightarrow \hat{X}$. If $\mathcal{E}_i \in I_{\text{inf}}((\hat{X}, M)/(\text{Spf}(V), N))^{lf}$ and there exists \mathcal{E}_c in the category $I_{\text{conv}}((U, M)/(\text{Spf}(V), N))^{lf}$ such that $\tilde{\Phi}(\mathcal{E}_c) = \mathcal{E}_i$, then $\mathcal{E}_i \in \tilde{\Phi}(I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))^{lf})$.*

Proof. Let us consider the sheaf

$$\mathcal{F} = z_{n*} \mathcal{H}om((K \otimes \mathcal{O}_{T_n(1)}) \otimes \mathcal{E}_i, \mathcal{E}_i \otimes (K \otimes \mathcal{O}_{T_n(1)})).$$

The following sequence is exact:

$$0 \rightarrow \mathcal{F} \xrightarrow{a} (\mathcal{F} \otimes \mathcal{O}_{\hat{X}(1)^{\text{an}}}) \oplus j_* j^*(\mathcal{F}) \xrightarrow{b} j_* j^*(\mathcal{F} \otimes \mathcal{O}_{\hat{X}(1)^{\text{an}}}),$$

because \mathcal{F} is a projective $z_{n*}(K \otimes \mathcal{O}_{T_n(1)})$ -module, and the following sequence is exact by proposition 5.2.8 of [Sh1] and [Og] lemma 2.14

$$0 \rightarrow z_{n*}(K \otimes \mathcal{O}_{T_n(1)}) \rightarrow \mathcal{O}_{\hat{X}(1)^{\text{an}}} \oplus j_* j^* z_{n*}(K \otimes \mathcal{O}_{T_n(1)}) \rightarrow j_* j^* \mathcal{O}_{\hat{X}(1)^{\text{an}}}.$$

Viewing \mathcal{E}_c as a convergent stratification we have a map

$$\epsilon'_n : j_* j^* z_{n*}((K \otimes \mathcal{O}_{T_n(1)}) \otimes \mathcal{E}_{i, \hat{X}}) \rightarrow j_* j^* z_{n*}(\mathcal{E}_{i, \hat{X}} \otimes (K \otimes \mathcal{O}_{T_n(1)}));$$

on the other hand the log infinitesimal isocrystal \mathcal{E}_i induces a stratification

$$\epsilon' : \mathcal{O}_{\hat{X}(1)^{\text{an}}} \otimes \mathcal{E}_{i, \hat{X}} \rightarrow \mathcal{E}_{i, \hat{X}} \otimes \mathcal{O}_{\hat{X}(1)^{\text{an}}}.$$

The pair (ϵ'_n, ϵ') lies in $\text{Ker}(b)$, then there exists an $\epsilon_n \in \mathcal{F}$ such that $a(\epsilon_n) = (\epsilon'_n, \epsilon')$ that defines a convergent stratification, i.e. a convergent isocrystal $\bar{\mathcal{E}}_c$. Moreover one can verify that $\tilde{\Phi}(\bar{\mathcal{E}}_c) = \mathcal{E}_i$. \square

Now we want to apply proposition 28 choosing as U the subset

$$\hat{X}_{\hat{f}-triv} = \{x \in \hat{X} \mid (\hat{f}^*N)_{\bar{x}} = (M)_{\bar{x}}\}.$$

Let us prove that it is open and dense in \hat{X} . Clearly $\hat{X}_{\hat{f}-triv}$ is homeomorphic to X_{k,f_k-triv} so it will be sufficient to prove that X_{k,f_k-triv} is open dense in X_k . But this follows from proposition 2.3.2 of [Sh1] because the special fiber is reduced.

Now applying proposition 28 we can restrict ourselves to the case in which $\hat{f}^*N = M$. As Shiho notices the hypothesis $\hat{f}^*N = M$ gives an equivalence of categories

$$I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N)) \cong I_{conv}((\hat{X}, \mathrm{triv})/(\mathrm{Spf}(V), \mathrm{triv})),$$

where with triv we indicate the trivial log structure and

$$I_{inf}((\hat{X}, M)/(\mathrm{Spf}(V), N)) \cong I_{inf}((\hat{X}, \mathrm{triv})/(\mathrm{Spf}(V), \mathrm{triv})).$$

So we reduce to the case of trivial log structures, as we wanted. We will characterize the essential image using certain type of stratification that we call special.

Definition 29. Let (E, ϵ_n) be an object of $\widehat{\mathrm{Str}}(\hat{X}/\mathrm{Spf}(V))$ and let \tilde{E} be a coherent p -torsion-free $\mathcal{O}_{\hat{X}}$ -module such that $K \otimes \tilde{E} = E$; we say that (E, ϵ_n) is special if there exists a sequence of integers $k(n)$ for $n \in \mathbb{N}$ such that:

(i) $k(n) = o(n)$ for $n \rightarrow \infty$,

(ii) the restriction of the map $p^{k(n)}\epsilon_n$ to $p_{2,n}^*(\tilde{E})$ has image contained in $p_{1,n}^*(\tilde{E})$ and the restriction of the map $p^{k(n)}\epsilon_n^{-1}$ to $p_{1,n}^*(\tilde{E})$ has image contained in $p_{2,n}^*(\tilde{E})$.

This definition is a small modification of definition of special stratification given by Shiho ([Sh1] Definition 5.2.12). Our definition of special is weaker than Shiho's definition: every special object in the sense of Shiho is special in our sense and allows us to characterize the essential image of the functor Φ .

Let us see now that the definition of special is well-posed.

Proposition 30. Definition 29 is independent of the choice of the p -torsion-free sheaf \tilde{E} .

Proof. Suppose that (E, ϵ_n) is special and that the conditions in definition 29 are verified for a given coherent p -torsionfree $\mathcal{O}_{\hat{X}}$ -module \tilde{E} such that $K \otimes \tilde{E} = E$. We take an other p -torsion-free $\mathcal{O}_{\hat{X}}$ -module \tilde{F} such that $K \otimes \tilde{F} = E$ and we want to prove that the same conditions are verified. From [Og] proposition 1.2 we know the following isomorphisms

$$K \otimes \mathrm{Hom}_{\mathcal{O}_{\hat{X}}}(\tilde{E}, \tilde{F}) \cong \mathrm{Hom}_{K \otimes \mathcal{O}_{\hat{X}}}(K \otimes \tilde{E}, K \otimes \tilde{F}) \cong \mathrm{End}_{K \otimes \mathcal{O}_{\hat{X}}}(E).$$

If we take the identity as endomorphism of E we know that there exists a power of p , say p^a , such that the multiplication by p^a is a morphism from \tilde{E} to \tilde{F} ; moreover this morphism is injective because \tilde{F} is p -torsion-free.

In the same way we can prove that there exists a b such that the multiplication by p^b is an injective morphism between \tilde{F} and \tilde{E} .

If we consider now the morphism $p^{k(n)+a+b}\epsilon_n$ then we have that the restriction of this to $p_{2,n}^*(\tilde{F})$ goes to $p_{1,n}^*(\tilde{F})$.

Arguing analogously for ϵ_n^{-1} we are done. \square

Let us see, first, that every object in the essential image of the functor Φ is special. Following Shiho [Sh1], proposition 3.2.14 and proposition 5.2.6, both in the case of trivial log structures, we have the equivalences of categories

$$I_{conv}(\hat{X}/\mathrm{Spf}(V)) = \mathrm{Str}'(\hat{X}/\mathrm{Spf}(V)),$$

$$I_{inf}(\hat{X}/\mathrm{Spf}(V)) = \widehat{Str}(\hat{X}/\mathrm{Spf}(V)).$$

The functor Φ induces the functor

$$\alpha : Str'(\hat{X}/\mathrm{Spf}(V)) \rightarrow \widehat{Str}(\hat{X}/\mathrm{Spf}(V)).$$

From now on we may work locally.

We are reduced to the situation where \hat{X} is formally smooth and we call dx_1, \dots, dx_l a basis of $\Omega_{\hat{X}/\mathrm{Spf}(V)}^1$. Let us call ξ_1, \dots, ξ_l the dual basis of dx_1, \dots, dx_l where $\xi_j = 1 \otimes x_j - x_j \otimes 1$; we will indicate (ξ_1, \dots, ξ_l) with ξ and an l -ple of natural numbers $(\beta_1, \dots, \beta_l)$ with β . We will use multi-index notations denoting $\prod_j \xi_j^{\beta_j}$ by ξ^β and $\beta_1 + \dots + \beta_l$ by $|\beta|$.

We call \hat{X}^n the n -th infinitesimal neighborhood of \hat{X} in $\hat{X} \times_{\mathrm{Spf} V} \hat{X}$. By a formal version of proposition 2.6 of [BeOg] we know that $\mathcal{O}_{\hat{X}^n}$ is a free $\mathcal{O}_{\hat{X}}$ -module generated by $\{\xi^\beta : |\beta| \leq n\}$, so that we can write $\mathcal{O}_{\hat{X}^n} = \mathcal{O}_{\hat{X}}[[\xi]]/(\xi^\beta, |\beta| = n+1)$.

We want to give a local description also for the universal system of enlargement $\{T_n\}_n$ of \hat{X} in $\hat{X} \times_{\mathrm{Spf} V} \hat{X}$. By [Og] remark 2.6.1 we know that T_n is isomorphic to the n -th universal enlargement of \hat{X} in $\hat{X} \times_{\mathrm{Spf} V} \hat{X}|_{\hat{X}}$, the formal completion of $\hat{X} \times_{\mathrm{Spf} V} \hat{X}$ along \hat{X} . Using this and the local description given in the proof of proposition 2.3 of [Og] we can write $\mathcal{O}_{T_n} = \mathcal{O}_{\hat{X}}\{\xi, \xi^\beta/p \mid |\beta| = n+1\}$. By universality of blowing up there exists a unique map ψ_n such that the following diagram is commutative

$$\begin{array}{ccc} & \hat{X}^n & \\ \psi_n \swarrow & \downarrow & \\ T_n & \xrightarrow{\quad} & \hat{X} \times_{\mathrm{Spf} V} \hat{X}. \end{array} \tag{11}$$

The functor α is induced by the pull back of ψ_n and in local coordinates is given by

$$\mathcal{O}_{T_n} = \mathcal{O}_{\hat{X}}\{\xi, \xi^\beta/p \mid (|\beta| = n+1)\} \rightarrow \mathcal{O}_{\hat{X}}[[\xi]]/(\xi^\beta, |\beta| = n+1).$$

Proposition 31. *If (E, ϵ_n) is in $\widehat{Str}(\hat{X}/\mathrm{Spf}(V))$ and it is in the image of the functor α , then it is special.*

Proof. Let (E, ϵ'_n) be an element of $Str'(\hat{X}/\mathrm{Spf}(V))$ such that $\alpha(E, \epsilon'_n) = (E, \epsilon_n)$, with

$$q_{2,n}^* E \xrightarrow{\epsilon'_n} q_{1,n}^* E,$$

where $q_{i,n}$ are the projections from T_n to \hat{X} , that exist by definition of convergent stratification.

We note that $q_{1,n}^* E = E \otimes_{\mathcal{O}_{\hat{X}}} \mathcal{O}_{\hat{X}}[[\xi]]\{\xi^\beta/p \mid (|\beta| = n+1)\}$ is embedded in $\prod_\beta E \xi^\beta$ and so we have a map, that we call ϵ' , which is the composition of

$$E \rightarrow q_{2,n}^* E \xrightarrow{\epsilon'_n} q_{1,n}^* E \rightarrow \prod_\beta E \xi^\beta.$$

Let us note that ϵ' does not depend from n , a consequence of the fact that the maps ϵ'_n coming from the convergent stratification are compatible. Using the isomorphisms ϵ_n that define the stratification (E, ϵ_n)

$$p_{2,n}^* E \xrightarrow{\epsilon_n} p_{1,n}^* E = \prod_{|\beta| \leq n} E \xi^\beta,$$

where $p_{i,n}$ are the projections $p_{i,n} : X^n \rightarrow X$, we can define a map

$$E \rightarrow p_{2,n}^* E \xrightarrow{\epsilon_n} p_{1,n}^* E = \prod_{|\beta| \leq n} E \xi^\beta. \tag{12}$$

The fact that $\alpha(E, \epsilon'_n) = (E, \epsilon_n)$ means that, if we call pr the projection

$$\prod_{\beta} E\xi^{\beta} \xrightarrow{pr} \prod_{|\beta| \leq n} E\xi^{\beta},$$

the map in (12) coincides with the map

$$E \xrightarrow{\epsilon'} \prod_{\beta} E\xi^{\beta} \xrightarrow{pr} \prod_{|\beta| \leq n} E\xi^{\beta}.$$

Let \tilde{E} be p -torsion-free $\mathcal{O}_{\hat{X}}$ -module such that $K \otimes \tilde{E} = E$; so $q_{1,n}^*(\tilde{E}) = \tilde{E} \otimes \mathcal{O}_{\hat{X}}[[\xi]]\{\xi^{\beta}/p \mid (|\beta| = n+1)\}$ and this is embedded in $\prod_{\beta} \tilde{E}\xi^{\beta}/p^{\lfloor \frac{|\beta|}{n+1} \rfloor}$.

Thus there exists a in \mathbb{N} such that

$$p^a \epsilon'(\tilde{E}) \subset \prod_{\beta} \tilde{E} \frac{\xi^{\beta}}{p^{\lfloor \frac{|\beta|}{n+1} \rfloor}}$$

and, if we call π_{β} the projection $\prod_{\beta} E\xi^{\beta} \rightarrow E$, then

$$p^{a+\lfloor \frac{|\beta|}{n+1} \rfloor} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subset \tilde{E};$$

therefore there exists a sequence $b_n(\beta)$ that tends to infinity when $|\beta|$ goes to infinity such that

$$p^{\lfloor \frac{|\beta|}{n} \rfloor} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subset p^{b_n(\beta)} \tilde{E}. \quad (13)$$

If we define now

$$a(k) := \min\{a \in \mathbb{N} \mid p^a \pi_{\beta} \circ \epsilon'(\tilde{E}) \subset \tilde{E} \text{ for all } \beta \text{ such that } |\beta| \leq k\},$$

then $p^{a(k)} \epsilon'_k(\tilde{E}) \subset \prod_{|\beta| \leq k} \tilde{E}\xi^{|\beta|}$, which means that $p^{a(k)} \epsilon'_k$ sends $p_{2,k}^*(\tilde{E})$ into $p_{1,k}^*(\tilde{E})$. So we are left to prove that $a(k) = o(k)$ for $k \rightarrow \infty$.

We notice, from the definition of $a(k)$, that $a(k)$ is an increasing sequence. If $a(k)$ is bounded, then we are done. Arguing by contradiction, then $a(k) \rightarrow \infty$ for $k \rightarrow \infty$. This means that there exists a sequence $\{k_i\}_i$ such that

$$0 < a(k_1) < a(k_2) < \dots < a(k_i) < a(k_{i+1}) < \dots$$

Then

$$p^{a(k_i)} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subseteq \tilde{E},$$

for every β such that $|\beta| \leq k_i$. Let us prove that

$$p^{a(k_i)} \pi_{\beta} \circ \epsilon'(\tilde{E}) \not\subseteq p\tilde{E} \quad (14)$$

for some β with $|\beta| = k_i$. Let us suppose that this is not true; this means that

$$p^{a(k_i)-1} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subseteq \tilde{E} \quad (15)$$

for every β such that $|\beta| = k_i$. Moreover for β with $|\beta| < k_i$ we have

$$p^{a(k_i)-1} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subseteq p^{a(k_{i-1})} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subseteq \tilde{E}.$$

Hence we have

$$p^{a(k_i)-1} \pi_{\beta} \circ \epsilon'(\tilde{E}) \subseteq \tilde{E}$$

for all β with $|\beta| \leq k_i$ and this contradicts the definition of $a(k_i)$'s, so (14) holds. If we now put together the formula (13) with $|\beta| = k_i$ and (14), we find that

$$\lim_{i \rightarrow \infty} \left(\left\lfloor \frac{k_i}{n} \right\rfloor - a(k_i) \right) = \infty,$$

so that there exists i_0 such that

$$0 \leq \frac{a(k_i)}{k_i} \leq \frac{1}{n}$$

for all $i \geq i_0$. Then for any $k \geq k_{i_0}$ we can find some k_i with $k_i \leq k \leq k_{i+1} - 1$ and then

$$0 \leq \frac{a(k)}{k} \leq \frac{a(k_i)}{k_i} \leq \frac{1}{n}.$$

Hence we have that $\limsup_k \frac{a(k)}{k} \leq \frac{1}{n}$. Since this is true for any n , we have that $a(k) = o(k)$. □

Now we want to prove the converse: that every special object is in the image of functor α . This is proven by Shiho in proposition 5.2.13 of [Sh1] for his special objects, but the proof works also in our case.

Proposition 32. *If (E, ϵ_n) is a special stratification on \hat{X} , then there exists $(E', \epsilon'_n) \in \text{Str}'(\hat{X}, \text{Spf}(V))$ such that $\alpha((E', \epsilon'_n)) = (E, \epsilon_n)$.*

So we have a complete characterization in term of stratification of the differential equations coming from a log-convergent isocrystal.

We want to describe the property of being special in term of radius of convergence. We will use the formalism given in [LS] in the local situation described before proposition 31. If we have an element $(E, \epsilon_n) \in \widehat{\text{Str}}(\hat{X}/\text{Spf}(V))$, then we can take the inverse limit of the map that we considered in the proof of the proposition 31

$$E \rightarrow p_{2,n}^* E \xrightarrow{\epsilon_n} p_{1,n}^* E = \prod_{|\beta| \leq n} E \xi^\beta,$$

and we obtain

$$\theta : E \rightarrow \varprojlim p_{2,n}^* E \xrightarrow{\varprojlim \epsilon_n} \varprojlim p_{1,n}^* E = \prod_{\beta} E \xi^\beta.$$

According to definition 4.4.1 of [LS] we can say that a section $s \in \Gamma(\hat{X}, E)$ is η -convergent, with $\eta < 1$ for the stratification (E, ϵ_n) if

$$\theta(s) \in \Gamma(\hat{X}, E \otimes_{\hat{X}} \{\frac{\xi}{\eta}\}).$$

Definition 33. *The radius of convergence of the section s for the stratification (E, ϵ_n) is defined as*

$$R(s) = \sup\{\eta \mid s \text{ is } \eta\text{-convergent}\}.$$

And the radius of convergence of the stratification (E, ϵ_n) is

$$R((E, \epsilon_n), \hat{X}) = \inf_{s \in \Gamma(\hat{X}, E)} R(s).$$

Proposition 34. *A stratification (E, ϵ_n) is special if and only if its radius of convergence is equal to 1.*

Proof. We know that $\widehat{Str}(\hat{X}/\mathrm{Spf}(V))$ is equivalent to the category $\widehat{\mathrm{MIC}}(\hat{X}/\mathrm{Spf}(V))$. By lemma 5.2.15 of [Sh1] we can write the map θ locally. Following the notation that we recalled before proposition 31 we denote by $\{D_\beta\}_{0 \leq |\beta| \leq n}$ the dual basis of $\{\xi^\beta\}_{0 \leq |\beta| \leq n}$ in $\mathrm{Diff}^n(\mathcal{O}_{\hat{X}}, \mathcal{O}_{\hat{X}})$, the differential operators of order $\leq n$ and in particular we indicate with $D_{(i)} := D_{(0, \dots, 1, \dots, 0)}$ with 1 at the i -th place.

With this notation

$$\theta(e) = \sum_{\beta} \frac{1}{\beta!} \nabla_{\beta}(e) \otimes \xi^{\beta}$$

with $\nabla_{\beta} := (\mathrm{id} \otimes D_{(1)} \circ \nabla)^{\beta_1} \circ \dots \circ (\mathrm{id} \otimes D_{(l)} \circ \nabla)^{\beta_l}$. Given an \tilde{E} , as in the definition 29, then the fact of being special can be translated as follows: there exists a sequence of integers $a(n)$ such that $a(n) = o(n)$ for $n \rightarrow \infty$ and that

$$p^{a(n)} \frac{\nabla_{\beta}(e)}{\beta!} \in \tilde{E}$$

for e in \tilde{E} and for any multi index β such that $|\beta| \leq n$. Let us see that the radius of convergence of a section $e \in \tilde{E}$ is 1, because $e \in \tilde{E}$ is η -convergent for every η , i.e. for every η

$$\theta(e) = \sum_{\beta} \frac{1}{\beta!} \nabla_{\beta}(e) \otimes \xi^{\beta} \in \Gamma(\hat{X}, E \otimes \mathcal{O}_{\hat{X}}\{\frac{\xi}{\eta}\}).$$

To prove it we have to show that if we denote by $\|\cdot\|$ the p -adic Banach norm on E such that $\|\tilde{E}\| = 1$

$$\|\frac{1}{\beta!} \nabla_{\beta}(e)\| \eta^{|\beta|} \rightarrow 0, \forall \eta < 1.$$

This is clearly true because, fixed an n , the following estimate holds:

$$\|\frac{1}{\beta!} \nabla_{\beta}(e)\| \eta^{|\beta|} = p^{a(n)} \|p^{a(n)} \frac{1}{\beta!} \nabla_{\beta}(e)\| \eta^{|\beta|} \leq p^{a(n)} \eta^{|\beta|}$$

for every β such that $|\beta| \leq n$, because $p^{a(n)} \frac{1}{\beta!} \nabla_{\beta}(e) \in \tilde{E}$.

This means that $\|\frac{1}{\beta!} \nabla_{\beta}(e)\| \eta^{|\beta|} \rightarrow 0$ since

$$0 \leq \|\frac{1}{\beta!} \nabla_{\beta}(e)\| \eta^{|\beta|} \leq p^{a(n)} \eta^{|\beta|}$$

and $p^{a(n)} \eta^{|\beta|} \rightarrow 0$ because $a(n) = o(n)$. So we can say that

$$R(e) = 1 \forall e \in \tilde{E},$$

and if we take $s \in E$, then there exists a positive integer k such that $p^k s \in \tilde{E}$, so that $R(p^k s) = 1$; moreover we know that s is η -convergent if and only if $p^k s$ is η convergent so

$$R(p^k s) = R(s)$$

and we can conclude that our stratification has radius of convergence 1.

The converse is also true: if (E, ϵ_n) is such that $R((E, \epsilon_n), \hat{X}) = 1$, then (E, ϵ_n) is special. We choose an \tilde{E} coherent $\mathcal{O}_{\hat{X}}$ -module p -torsion free such that $K \otimes \tilde{E} = E$; for every $e \in \tilde{E}$ let $a(n, e)$ the minimal integer such that

$$p^{a(n, e)} \frac{\nabla_{\beta}(e)}{\beta!} \in \tilde{E}$$

for any β with $|\beta| \leq n$. Since $R((E, \epsilon_n), \hat{X}) = 1$, then

$$\max_{|\beta| \leq n} \left(\left\| \frac{\nabla_{\beta}(e)}{\beta!} \right\| \right) \eta^n \leq \max_{|\beta| \leq n} \left(\left\| \frac{\nabla_{\beta}(e)}{\beta!} \right\| \eta^{\frac{|\beta|}{2}} \right) \eta^{\frac{n}{2}} \leq (\mathrm{const}) \eta^{\frac{n}{2}},$$

so that for $n \rightarrow \infty$ $p^{a(n,e)}\eta^n \rightarrow 0$ for any $\eta < 1$. This means that $a(n,e) = o(n)$ for any $e \in \tilde{E}$. Now if $e \in \tilde{E}$, then we can write $e = \sum_i^l f_i e_i$ where $f_i \in \mathcal{O}_{\hat{X}}$ and e_i 's are generator of \tilde{E} which is finitely generated $\mathcal{O}_{\hat{X}}$ module, and we put $a(n) := \max_{1 \leq i \leq l} a(n, e_i)$ (let us note that $a(n) = o(n)$). If we denote by d_{β} the operator ∇_{β} for the trivial stratification $(K \otimes \mathcal{O}_{\hat{X}}, \text{id})$, then for any $f \in \mathcal{O}_{\hat{X}}$ we have $\frac{d_{\beta}(f)}{\beta!} \in \mathcal{O}_{\hat{X}}$ for any β . Therefore, for any $e = \sum_i^l f_i e_i \in \tilde{E}$, we have

$$p^{a(n)} \frac{\nabla_{\beta}(e)}{\beta!} = \sum_{i=1}^l \sum_{0 \leq \gamma \leq \beta} \frac{d_{\gamma} f_i}{\gamma!} \left(p^{a(n)} \frac{\nabla_{\beta-\gamma} e_i}{(\beta-\gamma)!} \right) \in \tilde{E}.$$

Hence (E, ϵ_n) is special. \square

7 Description of the semistable case

In what follows we suppose that X is proper semistable variety over V , which means that locally for the étale topology there is an étale map

$$X \xrightarrow{\text{ét}} \text{Spec} \frac{V[x_1, \dots, x_n, y_1, \dots, y_m]}{x_1 \cdots x_r - \pi}.$$

We call M_{X_k} the log structure on X induced by the special fiber X_k that is a divisor with normal crossing, so locally for the étale topology it admits a chart given by

$$\mathbb{N}^r \rightarrow \frac{V[x_1, \dots, x_n, y_1, \dots, y_m]}{x_1 \cdots x_r - \pi}$$

that sends e_i to x_i , where $e_i = (0, \dots, 1, \dots, 0)$ with 1 at the i -th place.

We consider the log structure N induced by the closed point of $\text{Spec}(V)$ that has a chart given by

$$\mathbb{N} \rightarrow V,$$

that maps 1 to π . This is explained in [Ka] example 2.5 (1) and example 3.7 (2).

We also consider a normal crossing divisor D on X that locally for the étale topology is defined by the equation $\{y_1 \cdots y_s = 0\}$ and we indicate by M_D the log structure induced by D on X .

We consider on X the log structure $M = M_{X_k} \oplus M_D$, that corresponds to the log structure induced by the divisor with normal crossing $X_k \cup D$ in X (let us remark that with the notation $M_{X_k} \oplus M_D$ we indicate the sum in the category of log structures); the structural morphism extends to a log smooth morphism of log schemes $(X, M) \rightarrow (\text{Spec}(V), N)$. Moreover the special fiber is reduced, hence the hypothesis stated at the beginning of section 5 are satisfied.

If we denote by \hat{D} the p -adic completion of D , then we have a diagram

$$\begin{array}{ccc} D_k & \hookrightarrow & \hat{D} \\ \downarrow & & \downarrow \\ X_k & \hookrightarrow & \hat{X} \\ \downarrow & & \downarrow \\ \text{Spec}(k) & \hookrightarrow & \text{Spf}(V). \end{array}$$

We suppose that étale locally on \hat{X} we have the following diagram

$$\begin{array}{ccc} \hat{D} = \bigcup_{j=1}^s D_j & \hookrightarrow & \hat{X} \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y_j = 0\} & \hookrightarrow & \text{Spf}(V\{x_1, \dots, x_n, y_1, \dots, y_m\}/(x_1 \cdots x_r - \pi)) \end{array} \quad (16)$$

which is cartesian with the vertical maps that are étale and the horizontal maps closed immersions. If \hat{X}_{sing} and \hat{D}_{sing} are the singular loci of \hat{X} and \hat{D} respectively, then we will use the following notations:

$$\begin{aligned}\hat{X}^\circ &= \hat{X} - (\hat{X}_{sing} \cup \hat{D}_{sing}) \\ \hat{D}^\circ &= \hat{D} - (\hat{X}_{sing} \cup \hat{D}_{sing}) = \hat{X}^\circ \cap \hat{D}.\end{aligned}$$

When we consider the situation étale locally and fix a diagram (16), we have a decomposition of the formal schemes $\hat{X} - \hat{X}_{sing}$, \hat{X}° and \hat{D}° which will be useful later. First let \hat{X}_i° be the open formal scheme of \hat{X} defined by pullback of the open formal scheme of $\mathrm{Spf}(V\{x_1, \dots, x_n, y_1, \dots, y_m\}/(x_1 \cdots x_r - \pi))$ on which all the $x_{i'}$'s for $i' \neq i$ are invertible and let $\hat{X}_{i,j}^\circ$ be the open formal subscheme of \hat{X}_i° defined by étale pullback of the open formal subscheme of $\mathrm{Spf}(V\{x_1, \dots, x_n, y_1, \dots, y_m\}/(x_1 \cdots x_r - \pi))$, where $x_{i'}$ are invertible $\forall i' \neq i, 1 \leq i \leq r$ and $y_{j'}$ are invertible $\forall j' \neq j, 1 \leq j \leq s$.

Moreover we will indicate with $\hat{D}_{i,j}^\circ$ the set $\hat{X}_{i,j}^\circ \cap \hat{D} = \hat{X}_{i,j}^\circ \cap \hat{D}_j$, that is the open formal subscheme of \hat{D}_j defined by pullback of the open formal subscheme of $\mathrm{Spf}(V\{x_1, \dots, x_n, y_1, \dots, \hat{y}_j, \dots, y_m\}/(x_1 \cdots x_r - \pi))$, where all the $x_{i'}$ and the $y_{j'}$ are invertible for all $i' \neq i, 1 \leq i \leq r, \forall j' \neq j, 1 \leq j \leq s$. In the previous line \hat{y}_j in $\mathrm{Spf}(V\{x_1, \dots, x_n, y_1, \dots, \hat{y}_j, \dots, y_m\}/(x_1 \cdots x_r - \pi))$ means that the coordinate y_j is missing.

With this notations we have the following relations:

$$\begin{aligned}\coprod_i \hat{X}_i^\circ &= \hat{X} - \hat{X}_{sing}, & \coprod_{i,j} \hat{X}_{i,j}^\circ &= \hat{X}^\circ, \\ \coprod_{i,j} \hat{D}_{i,j}^\circ &= \hat{D}^\circ.\end{aligned}$$

Note that this decomposition is defined only if we work étale locally and we fix a diagram as (16). If we denote by the subscript K the rigid analytic space associate to a formal scheme, then the sets $\hat{D}_{i,j;K}^\circ$ and $\hat{X}_{i,j;K}^\circ$ can be described as follows:

$$\begin{aligned}\hat{X}_{i,j;K}^\circ &= \{p \in \hat{X}_K \mid \forall i' \neq i \ |x_{i'}(p)| = 1, \forall j' \neq j \ |y_{j'}(p)| = 1\}, \\ \hat{D}_{i,j;K}^\circ &= \{p \in \hat{X}_K \mid \forall i' \neq i \ |x_{i'}(p)| = 1, \forall j' \neq j \ |y_{j'}(p)| = 1, y_j(p) = 0\}.\end{aligned}$$

Finally we will denote by \hat{U} the open formal subscheme complement of \hat{D} in \hat{X} .

8 Log convergent isocrystals with exponents in Σ

We consider now the category of locally free log convergent isocrystals on \hat{X} , that we denote, as before, by $I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$. By remark 5.1.3 of [Sh1] we know that there is an equivalence of categories between $((X_k, M)/(\mathrm{Spf}(V), N))_{conv}$, the log convergent site on the special fiber, and $((\hat{X}, M)/(\mathrm{Spf}(V), N))_{conv}$, the log convergent site on the lifting, hence an equivalence of categories between $I_{conv}((\hat{X}, M)/(\mathrm{Spf}(V), N))^{lf}$ and $I_{conv}((X_k, M)/(\mathrm{Spf}(V), N))^{lf}$. As we saw in section 5, through the functor Φ we can associate to a locally free convergent log isocrystal \mathcal{E} on \hat{X} a locally free infinitesimal log isocrystal $\tilde{\Phi}(\mathcal{E})$. Using the terminology of [Ke] and [Sh6], in the local situation as in (16), $\tilde{\Phi}(\mathcal{E})$ induces a log- ∇ -module E on \hat{X}_K with respect to y_1, \dots, y_s , that means a locally free coherent module E on \hat{X}_K and an integrable connection

$$\nabla : E \rightarrow E \otimes \omega_{\hat{X}_K/K}^1,$$

where $\omega_{\hat{X}_K/K}^1$ is the coherent sheaf on \hat{X}_K associated to the isocoherent sheaf $K \otimes \omega_{(\hat{X}, M)/(\mathrm{Spf}(V), N)}^1$ on \hat{X} . If we are in the situation of (16) we can write $\omega_{\hat{X}_K/K}^1$ more explicitly: if we denote by $\Omega_{\hat{X}_K/K}^1$ the sheaf of continuous classical 1-differentials on the rigid analytic space \hat{X}_K , then

$$\omega_{\hat{X}_K/K}^1 = (\Omega_{\hat{X}_K/K}^1 / K \oplus \bigoplus_{j=1}^s \mathcal{O}_{\hat{X}_K} d\log y_j) / L,$$

where L is the coherent sub $\mathcal{O}_{\hat{X}_K}$ generated by $(dy_j, 0) - (0, y_j d\log y_j)$ for $1 \leq j \leq s$. Fixed a $j' \in \{1, \dots, s\}$, i.e a component $\hat{D}_{j';K} = \{y'_{j'} = 0\}$ of \hat{D}_K , then there is a natural immersion of

$$\Omega^1_{\hat{X}_K}/K \oplus \bigoplus_{j \neq j'} \mathcal{O}_{\hat{X}_K} d\log y_j \rightarrow \omega^1_{\hat{X}_K/K}$$

and we call $M_{j'}$ the image. The endomorphism $\text{res}_j \in \text{End}_{\mathcal{O}_{\hat{D}_{j;K}}}(E \otimes_{\mathcal{O}_{\hat{X}_K}} \mathcal{O}_{\hat{D}_{j;K}})$ obtained tensoring by $\mathcal{O}_{\hat{D}_{j;K}}$ the following map

$$E \rightarrow E \otimes \omega^1_{\hat{X}_K/K} \rightarrow E \otimes \omega^1_{\hat{X}_K/K}/M_j$$

is called the residue of E along $\hat{D}_{j;K}$. Thanks to proposition 1.5.3 of [BaCh] we know that there exists a minimal and monic polynomial $P_j \in K[T]$ such that $P_j(\text{res}_j) = 0$. The exponents of (E, ∇) along $\hat{D}_{j;K}$ are the roots of P_j .

We fix a set $\Sigma = \prod_{h=1}^k \Sigma_h \in \mathbb{Z}_p^k$, where k is the number of the irreducible components of $\hat{D} = \bigcup_{h=1}^k \hat{D}^h$ in \hat{X} .

If there exists an étale covering $\coprod_l \phi_l : \coprod_l \hat{X}_l \rightarrow \hat{X}$ such that every \hat{X}_l has a diagram as in (16), then we can define a function of sets $h : \{1, \dots, r\} \times \{1, \dots, s\} \rightarrow \{1, \dots, k\}$ as follows: with the notation as in the previous paragraph $\phi_l(\hat{D}_{i,j,l}^\circ)$ is contained in one irreducible component of \hat{D} , which we denote by $\hat{D}^{h(i,j)}$. We denote by $\Sigma_{h(i,j)}$ the factor of Σ corresponding to the component $\hat{D}^{h(i,j)}$.

Definition 35. A locally free convergent isocrystal \mathcal{E} has exponents along \hat{D}_K in Σ if there exists an étale covering $\coprod_l \phi_l : \coprod_l \hat{X}_l \rightarrow \hat{X}$ such that every \hat{X}_l has a diagram

$$\begin{array}{ccc} \hat{D}_l = \bigcup_{j=1}^s \hat{D}_{j,l} & \hookrightarrow & \hat{X}_l \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y_{l,j} = 0\} & \hookrightarrow & \text{Spf} V\{x_{l,1}, \dots, x_{l,n}, y_{l,1}, \dots, y_{l,m}\} / (x_{l,1} \dots x_{l,r} - \pi) \end{array} \quad (17)$$

as in (16) with $\hat{D}_l := \phi_l^{-1}(\hat{D})$ such that for every l the log- ∇ -module E_l on $\hat{X}_{l;K}$ induced by \mathcal{E} has exponents along $\hat{D}_{j,l;K}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$, if $\phi_l(\hat{D}_{i,j,l}^\circ) \subset \hat{D}^{h(i,j)}$.

We will denote by $I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))^\Sigma$ or $I_{\text{conv}}(\hat{X}/\text{Spf}(V))^{\log, \Sigma}$ the category of locally free log convergent isocrystals with exponents in Σ .

In the next lemma we prove that the definition of isocrystals with exponents along \hat{D}_K in Σ is well posed.

Lemma 36. The notion of locally free log convergent isocrystal with exponents in Σ is independent of the choice of the étale covering and the diagram as in (16), which are chosen in definition 35.

Proof. Let us suppose that \mathcal{E} is a log convergent isocrystal with exponents along \hat{D}_K in Σ . It is sufficient to prove that for any étale morphism $\phi : \hat{X}' \rightarrow \hat{X}$, such that for \hat{X}' there exists a diagram

$$\begin{array}{ccc} \hat{D}' = \bigcup_{j'=1}^{s'} \hat{D}'_{j'} & \hookrightarrow & \hat{X}' \\ \downarrow & & \downarrow \\ \bigcup_{j'=1}^{s'} \{y'_{j'} = 0\} & \hookrightarrow & \text{Spf} V\{x'_1, \dots, x'_{n'}, y'_{1'}, \dots, y'_{m'}\} / (x'_1 \dots x'_{r'} - \pi) \end{array} \quad (18)$$

as in (16), with $\hat{D}' := \phi^{-1}(\hat{D})$, the log- ∇ -module E' on \hat{X}'_K induced by \mathcal{E} has exponents along $\hat{D}'_{j';K}$ in $\cap_{i'=1}^{r'} \Sigma_{h(i',j')}$, if $\phi(\hat{D}'_{i',j'}^\circ) \subset \hat{D}^{h(i',j')}$.

By hypothesis \mathcal{E} has exponents along \hat{D}_K in Σ , hence there exists an étale covering $\coprod_l \phi_l : \coprod_l \hat{X}_l \rightarrow \hat{X}$ such that every \hat{X}_l has a diagram

$$\begin{array}{ccc} \hat{D}_l = \bigcup_{j=1}^s \hat{D}_{j,l} & \hookrightarrow & \hat{X}_l \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y_{l,j} = 0\} & \hookrightarrow & \mathrm{Spf}V\{x_{l,1}, \dots, x_{l,n}, y_{l,1}, \dots, y_{l,m}\} / (x_{l,1} \dots x_{l,r} - \pi) \end{array} \quad (19)$$

as in (16) with $\hat{D}_l := \phi_l^{-1}(\hat{D})$ such that for every l the log- ∇ -module E_l on $\hat{X}_{l,K}$ induced by \mathcal{E} has exponents along $\hat{D}_{l,j;K}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$, if $\phi_l(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

Let us denote by \hat{X}'_l the fiber product $\hat{X}' \times_{\hat{X}} \hat{X}_l$ and by \hat{D}'_l , $\hat{D}'_{l,j'}$, $\hat{D}''_{j,l}$ the inverse image of \hat{D}_l , $\hat{D}'_{j'}$, $\hat{D}_{j,l}$ on \hat{X}'_l respectively. With this notation we have two diagrams on \hat{X}'_l :

$$\begin{array}{ccc} \hat{D}'_l = \bigcup_{j'=1}^{s'} \hat{D}'_{j',l} & \hookrightarrow & \hat{X}'_l \\ \downarrow & & \downarrow \\ \bigcup_{j'=1}^{s'} \{y'_{j'} = 0\} & \hookrightarrow & \mathrm{Spf}V\{x'_1, \dots, x'_{n'}, y'_1, \dots, y'_{m'}\} / (x'_1 \dots x'_{r'} - \pi) \end{array} \quad (20)$$

and

$$\begin{array}{ccc} \hat{D}'_l = \bigcup_{j=1}^s \hat{D}''_{j,l} & \hookrightarrow & \hat{X}'_l \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y_{j,l} = 0\} & \hookrightarrow & \mathrm{Spf}V\{x_{1,l}, \dots, x_{n,l}, y_{1,l}, \dots, y_{m,l}\} / (x_{1,l} \dots x_{r,l} - \pi). \end{array} \quad (21)$$

The diagram (20) is induced by (18) through p_2 , the projection on the second factor $\mathrm{pr}_2 : \hat{X}' \times_{\hat{X}} \hat{X}_l \rightarrow \hat{X}_l$; and the diagram (21) is induced by (19) through p_1 , the projection on the first factor $\mathrm{pr}_2 : \hat{X}' \times_{\hat{X}} \hat{X}_l \rightarrow \hat{X}'$. The log- ∇ -module E'_l induced by \mathcal{E} on $\hat{X}'_{l,K}$ has exponents along $\hat{D}''_{j,l;K}$ which are contained in the set of exponents of E_l along $\hat{D}_{j,l;K}$. This happens because the residue of E'_l along $\hat{D}''_{j,l;K}$, denoted by $\mathrm{res}''_{j,l}$ is the image of the residue of E_l along $\hat{D}_{j,l;K}$, denoted by $\mathrm{res}_{j,l}$, via the map

$$\mathrm{End}_{\mathcal{O}_{\hat{D}_{j,l;K}}} (E_l|_{\hat{D}_{j,l;K}}) \rightarrow \mathrm{End}_{\mathcal{O}_{\hat{D}''_{j,l;K}}} (E'_l|_{\hat{D}''_{j,l;K}}),$$

which is induced by the projection pr_2 . If $P_{j,l}$ is the minimal and monic polynomial such that $P_{j,l}(\mathrm{res}_{j,l}) = 0$, then $P_{j,l}(\mathrm{res}''_{j,l}) = 0$, so if we denoted by $P''_{j,l}$ the minimal and monic polynomial such that $P''_{j,l}(\mathrm{res}''_{j,l}) = 0$, then $P''_{j,l} \mid P_{j,l}$. So the roots of $P''_{j,l}$ are contained in the roots of $P_{j,l}$, which means that the exponents of E'_l along $\hat{D}''_{j,l}$ are a subset of the set of exponents of E_l along $\hat{D}_{j,l;K}$. Since for every (i,j) such that $\phi_l \circ \mathrm{pr}_2(\hat{D}''_{i,j,l})$ is contained in $\hat{D}^{h(i,j)}$ also $\phi_l(\hat{D}^\circ_{i,j,l})$ is contained in $\hat{D}^{h(i,j)}$ and viceversa, then we proved that for every (i,j) such that $\phi_l \circ \mathrm{pr}_2(\hat{D}''_{i,j,l})$ is contained in $\hat{D}^{h(i,j)}$ the exponents along $\hat{D}''_{j,l;K}$ are contained in $\cap_{i=1}^r \Sigma_{h(i,j)}$.

Now we want to look at the exponents of E'_l along $\hat{D}'_{j',l;K}$.

Let us put $\hat{X}'_l{}^\circ := \hat{X}'_l \cap \hat{X}^\circ$, $\hat{D}'_l{}^\circ := \hat{D}'_l \cap \hat{X}^\circ$, $\hat{D}'_{l,j'}{}^\circ := \hat{D}'_{l,j'} \cap \hat{X}^\circ$ and $\hat{D}''_{l,j}{}^\circ := \hat{D}''_{l,j} \cap \hat{X}^\circ$. Since the map

$$\mathrm{End}_{\mathcal{O}_{\hat{D}'_{l,j';K}}} (E'_l|_{\hat{D}'_{l,j';K}}) \rightarrow \mathrm{End}_{\mathcal{O}_{\hat{D}'_{l,j';K}}} (E'_l|_{\hat{D}'_{l,j';K}{}^\circ})$$

is injective it is enough to look at the exponents of $E'_l|_{\hat{X}'_l{}^\circ}$ along $\hat{D}'_{l,j'}{}^\circ$.

Let us note that $\hat{D}_l'^\circ$ is a relative normal crossing divisor in a smooth formal V -scheme; if $\hat{D}_l'^\circ = \bigcup_t \hat{C}_{t,l}$ is the decomposition of $\hat{D}_l'^\circ$ in irreducible components, from [NaSh] proposition A.0.3 and proposition A.0.7, we can deduce that $\hat{D}_{i',j',l}'^\circ$ and $\hat{D}_{i,j,l}''^\circ$, that are irreducible components of $\hat{D}_l'^\circ$, correspond to some $\hat{C}_{t,l}$'s. Thanks to what we have proven before we know that $E_l' |_{\hat{X}_{l,K}'^\circ}$ has exponents along $\hat{D}_{l,j,K}''^\circ$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$ where (i,j) are such that $\phi_l \circ \text{pr}_2(\hat{D}_{i,j,l}'^\circ)$ is contained in $\hat{D}^{h(i,j)}$. We now consider $\hat{D}_{j',l}'^\circ$, then $\hat{D}_{i',j',l}'^\circ$ will coincide with some $C_{t,l}$'s, so will correspond to some $\hat{D}_{i,j,l}''^\circ$. If $C_{t,l}$ is such that $\phi \circ \text{pr}_1(C_{t,l}) \subset \hat{D}^h$ then also $\phi_l \circ \text{pr}_2(C_{t,l}) \subset \hat{D}^h$ for the commutativity of the following diagram

$$\begin{array}{ccc} \hat{X}' \times_{\hat{X}} \hat{X}_l & \xrightarrow{\text{pr}_2} & \hat{X}_l \\ \downarrow \text{pr}_1 & & \downarrow \phi_l \\ \hat{X}' & \xrightarrow{\phi} & \hat{X} \end{array} \quad (22)$$

So we can conclude that the exponents of E_l' along $\hat{D}_{j',l,K}'^\circ$ are contained in $\cap_{i'=1}^{r'} \Sigma^{h(i',j')}$ with (i',j') such that $\phi \circ \text{pr}_1(\hat{D}_{i',j',l}'^\circ) \subset \hat{D}^{h(i',j')}$.

Finally we prove that E' has exponents along $\hat{D}_{j',K}'^\circ$ in $\cap_{i'=1}^{r'} \Sigma_{h(i',j')}$ with (i',j') such that $\phi(\hat{D}_{i',j',l}'^\circ) \subset \hat{D}^{h(i',j')}$. Having a surjective étale map $\coprod_l \hat{X}_l' \rightarrow \hat{X}'$, the thesis is reduced to prove that the induced map

$$\text{End}_{\mathcal{O}_{\hat{D}_{j',K}'^\circ}}(E' \otimes \mathcal{O}_{\hat{D}_{j',K}'^\circ}) \rightarrow \text{End}_{\mathcal{O}_{\coprod_l \hat{D}_{j',l,K}'^\circ}}(\coprod_l E_l' \otimes \mathcal{O}_{\coprod_l \hat{D}_{j',l,K}'^\circ}) \quad (23)$$

is injective.

If $\phi(\hat{D}_{i',j',l}'^\circ) \subset \hat{D}^{h(i',j')}$, then $\phi \circ \text{pr}_1(\hat{D}_{i',j',l}'^\circ) \subset \hat{D}^{h(i',j')}$ for every l and $\phi \circ \text{pr}_1(\coprod_l \hat{D}_{i',j',l}'^\circ)$ is contained in $\hat{D}^{h(i',j')}$. One can see that the residue of E' along $\hat{D}_{j',K}'^\circ$ goes via the map in (23) into the residue of $\coprod_l E_l'$ along $\coprod_l \hat{D}_{j',l,K}'^\circ$.

$$\Gamma(\hat{D}_{j',K}'^\circ, \mathcal{O}_{\hat{D}_{j',K}'^\circ}) \rightarrow \Gamma(\coprod_l \hat{D}_{j',l,K}'^\circ, \mathcal{O}_{\coprod_l \hat{D}_{j',l,K}'^\circ})$$

is injective, since $\coprod_l \hat{X}_l' \rightarrow \hat{X}'$ is étale surjective and then faithfully flat. \square

9 Log- ∇ -modules on polyannuli

We recall in this section the notion of log- ∇ -modules on some particular rigid space defined and used by Kedlaya in [Ke] and by Shiho in [Sh6].

An aligned interval is a interval I contained in $[0, \infty)$ such that any end point is contained in Γ^* with Γ^* the divisible closure of the image of the absolute value $|\cdot| : K^* \rightarrow \mathbb{R}^+$. An aligned interval is said to be quasi open if is open at any non zero end point. For an aligned interval we define a polyannulus as the rigid analytic space $A_K^n(I) = \{(t_1, \dots, t_n) \in \mathbb{A}_K^{n,rig} \mid |t_i| \in I \forall i = 1, \dots, n\}$.

If Y is a smooth rigid analytic space and y_1, \dots, y_s are global sections such that they are smooth and meet transversally, then for a subset $\Sigma = \prod_{j=1}^s \Sigma_j \subset \bar{K}^s$ we denote by $\text{LNM}_{Y,\Sigma}$ the category of log- ∇ -module on Y such that all the exponents along $\{y_j = 0\}$ are contained in Σ_j for every $j = 1, \dots, s$.

If Y is a smooth rigid analytic space and y_1, \dots, y_s are global sections such that they are smooth and meet transversally, then we set $\omega_{Y \times A_K^n([0,0])/K} = \omega_{Y/K}^1 \oplus \bigoplus_{i=1}^n \mathcal{O}_Y d \log t_i$. We define a log- ∇ -module (E, ∇) on $Y \times A_K^n([0,0])/K$ with respect to $y_1, \dots, y_s, t_1, \dots, t_n$ as a log- ∇ -module (E, ∇) on Y with respect to y_1, \dots, y_s with n commuting endomorphisms $\partial_i = t_i \frac{\partial}{\partial t_i}$ of (E, ∇) for $i = 1, \dots, n$. If we fix $\Sigma = \prod_{j=1}^s \Sigma_j \times \prod_{i=1}^n \Sigma_i \subset \bar{K}^{s+n}$, we can define a log- ∇ -module (E, ∇) with respect to $y_1, \dots, y_s, t_1, \dots, t_n$ on $Y \times A_K^n([0,0])/K$ with exponents in Σ if the log- ∇ -module (E, ∇) on Y has exponents along $\{y_j = 0\}$ in Σ_j and if the commuting endomorphisms $\partial_i = t_i \frac{\partial}{\partial t_i}$ have eigenvalues in Σ_i for every $i = 1, \dots, n$. Following Shiho we denote the

category of locally free log- ∇ -modules on $Y \times A_K^n([0,0])/K$ with exponents in Σ by $\text{LNM}_{Y \times A_K^n([0,0]), \Sigma}$. If I is an aligned interval and $\xi := (\xi_1, \dots, \xi_n) \in \bar{K}^n$, the log- ∇ -module denoted by (M_ξ, ∇_{M_ξ}) is the log- ∇ -module on $A_K^n(I)$ given by $(\mathcal{O}_{A_K^n(I)}, d + \sum_{j=1}^n \xi_j d \log t_j)$. We will define now the notion of Σ -unipotence for log- ∇ -modules on a product of a smooth rigid analytic space and a polyannulus ([Sh6] definition 1.3).

Definition 37. Let Y be a smooth rigid analytic space, y_1, \dots, y_s global sections whose zero loci are smooth and meet transversally, I an aligned interval and $\Sigma = \prod_{j=1}^{s+n} \Sigma_j \subset \bar{K}^{s+n}$. We say that $\mathcal{E} = (E, \nabla) \in \text{LNM}_{Y \times A_K^n(I)}$ is Σ -unipotent if after some finite extension of K there exists a filtration

$$0 \subset \mathcal{E}_1 \subset \dots \subset \mathcal{E}_n = \mathcal{E}$$

of subobjects such that every successive quotient $\mathcal{E}_i/\mathcal{E}_{i-1} = \pi_1^* \mathcal{F} \times \pi_2^*(M_\xi, \nabla_{M_\xi})$, where by π_1 we denote the first projection $\pi_1 : Y \times A_K^n(I) \rightarrow Y$, by π_2 the second projection $\pi_2 : Y \times A_K^n(I) \rightarrow A_K^n(I)$, by \mathcal{F} a log- ∇ -module $\in \text{LNM}_{X, \prod_{j=1}^s \Sigma_j}$ and by (M_ξ, ∇_{M_ξ}) the log- ∇ -module we defined before with $\xi \in \prod_{j=s+1}^{s+n} \Sigma_j$. We will denote by $\text{ULNM}_{Y \times A_K^n(I), \Sigma}$ the categories of Σ -unipotent log- ∇ -modules on $Y \times A_K^n(I)$.

Shiho ([Sh6] Definition 1.5) defines a functor

$$\mathcal{U}_I : \text{LNM}_{Y \times A_K^n([0,0]), \Sigma} \rightarrow \text{LNM}_{Y \times A_K^n(I), \Sigma}$$

that associates to a log- ∇ -module \mathcal{E} with respect to y_1, \dots, y_s on Y with n commuting endomorphisms N_i for $i = 1, \dots, n$ with eigenvalues on Σ_i a log- ∇ -module $\mathcal{U}_I(\mathcal{E})$ defined as the sheaf $\pi_1^* \mathcal{E}$ and the connection

$$v \mapsto \pi_1^*(\nabla)v + \sum_{i=1}^n \pi_1^*(N_i)(v) \otimes \frac{dt_i}{t_i}.$$

We recall here the definition of a non Liouville number, which we will use in the sequel.

Definition 38. An element α in \bar{K} is said to be p -adically non-Liouville if both the power series $\sum_{n \neq \alpha} \frac{x^n}{\alpha - n}$ and $\sum_{n \neq \alpha} \frac{x^n}{n - \alpha}$ have radius of convergence equal to 1.

As in definition 1.8 of [Sh6] we can define the following.

Definition 39. A set $\Sigma \subset \bar{K}$ is called (NID) (resp. (NLD)) if for any $\alpha, \beta \in \Sigma$, $\alpha - \beta$ is a non zero integer (resp. is p -adically non-Liouville). A set $\Sigma = \prod_{j=1}^s \Sigma_j \subset \bar{K}^s$ is called (NID) (resp. (NLD)) if for any $j = 1, \dots, s$ Σ_j is (NID) (resp. (NLD)).

We will use the following result ([Ke] 3.3.4, [Ke] 3.3.6, [Sh6] corollary 1.15 and [Sh6] corollary 1.16)

Theorem 40. Let Y be as before, I a quasi open interval and $\Sigma = \prod_{j=1}^{s+n} \Sigma_j \subset \bar{K}^{s+n}$ which is (NID) and (NLD) then the restriction of the functor \mathcal{U}_I to the Σ -unipotent log- ∇ -modules

$$\mathcal{U}_I : \text{ULNM}_{Y \times A_K^n([0,0]), \Sigma} \rightarrow \text{ULNM}_{Y \times A_K^n(I), \Sigma}$$

is an equivalence of categories. If I is an interval of positive length, then the functor \mathcal{U}_I is fully faithful.

10 Log overconvergent isocrystals

Before defining the category of log overconvergent isocrystals, we recall the notion of log tubular neighborhood given by Shiho in [Sh2] definition 2.2.5 with some restrictive hypothesis and in [Sh3] paragraph 2 in full generality. This is the log version of the tubular neighborhood defined by Berthelot in [Be].

Given a closed immersion of fine log formal schemes $i : (Z, M_Z) \hookrightarrow (\mathcal{Z}, M_{\mathcal{Z}})$, there exists a fine log formal scheme $(\mathcal{Z}^{ex}, M_{\mathcal{Z}^{ex}})$ and an associated homeomorphic closed exact immersion $i^{ex} : (Z, M_Z) \hookrightarrow (\mathcal{Z}^{ex}, M_{\mathcal{Z}^{ex}})$

such that the functor that associates i^{ex} to i is a right adjoint functor to the inclusion functor from the category of homeomorphic closed immersions of log formal schemes in the category of closed immersions of log formal schemes. The functor $i \mapsto i^{ex}$ is called the exactification functor and its existence is proven in [Sh3] proposition-definition 2.10.

Definition 41. Let $(Z, M_Z) \hookrightarrow (\mathcal{Z}, M_{\mathcal{Z}})$ be a closed immersion of log formal schemes, then the log tubular neighborhood $]Z[_{\mathcal{Z}}^{log}$ of (Z, M_Z) in $(\mathcal{Z}, M_{\mathcal{Z}})$ is defined as the rigid analytic space \mathcal{Z}_K^{ex} associated to the formal scheme \mathcal{Z}^{ex} . We can define the specialization map

$$sp :]Z[_{\mathcal{Z}}^{log} \rightarrow \hat{\mathcal{Z}},$$

where $\hat{\mathcal{Z}}$ is the completion of \mathcal{Z} along Z , as the composition of the usual specialization map $]Z[_{\mathcal{Z}}^{log} = \mathcal{Z}_K^{ex} \rightarrow \mathcal{Z}^{ex}$ with the map $\mathcal{Z}^{ex} \rightarrow \hat{\mathcal{Z}}$ induced by the morphism $\mathcal{Z}^{ex} \rightarrow \mathcal{Z}$.

We can notice that, if the closed immersion $i : (Z, M_Z) \hookrightarrow (\mathcal{Z}, M_{\mathcal{Z}})$ is exact, then $\mathcal{Z}^{ex} = \mathcal{Z}_K$ and the log tubular neighborhood $]Z[_{\mathcal{Z}}^{log}$ coincides with the classical tubular neighborhood.

We define the category of log overconvergent isocrystals for log pairs. Log pairs are defined in paragraph 4 of [Sh4] and in 2.1 of [ChTs] in the case of trivial log structures. A log pair is a pair $((X, M_X), (\bar{X}, M_{\bar{X}}))$ of fine log schemes in characteristic p endowed with a strict open immersion $(X, M_X) \hookrightarrow (\bar{X}, M_{\bar{X}})$. A morphism of log pairs $f : ((X, M_X), (\bar{X}, M_{\bar{X}})) \rightarrow ((Y, M_Y), (\bar{Y}, M_{\bar{Y}}))$ is a morphism of log schemes $f : \bar{X} \rightarrow \bar{Y}$ that verifies $f(X) \subset Y$. A log pair $((X, M_X), (\bar{X}, M_{\bar{X}}))$ over a log pair $((S, M_S), (\bar{S}, M_{\bar{S}}))$ is a log pair endowed with the structural morphism $f : ((X, M_X), (\bar{X}, M_{\bar{X}})) \rightarrow ((S, M_S), (\bar{S}, M_{\bar{S}}))$. We assume that all log pairs are log pairs over a given log pair $((S, M_S), (\bar{S}, M_{\bar{S}}))$. In paragraph 4 of [Sh4] there is a definition of log overconvergent isocrystals for log pairs over a log pair $((S, M_S), (\bar{S}, M_{\bar{S}}))$ with M_S isomorphic to the trivial log structure; we will give analogous definition in the case of non necessarily trivial M_S .

A log triple is a triple $((X, M_X), (\bar{X}, M_{\bar{X}}), (\mathcal{P}, M_{\mathcal{P}}))$ which consists of a log pair $((X, M_X), (\bar{X}, M_{\bar{X}}))$ and a log formal scheme $(\mathcal{P}, M_{\mathcal{P}})$ over a log formal scheme $(\mathcal{S}, M_{\mathcal{S}})$ endowed with a closed immersion $(\bar{X}, M_{\bar{X}}) \hookrightarrow (\mathcal{P}, M_{\mathcal{P}})$. Morphisms of log triples are defined in the natural way, as well as a log triple over another log triple. We will work only with triples $((X, M_X), (\bar{X}, M_{\bar{X}}), (\mathcal{P}, M_{\mathcal{P}}))$ over a fixed log triple $((S, M_S), (\bar{S}, M_{\bar{S}}), (\mathcal{S}, M_{\mathcal{S}}))$.

As in the classical case, for a log triple $((X, M_X), (\bar{X}, M_{\bar{X}}), (\mathcal{P}, M_{\mathcal{P}}))$ we can define a strict neighborhood W of $]X[_{\mathcal{P}}^{log}$ in $]\bar{X}[_{\mathcal{P}}^{log}$ to be an admissible open of $]\bar{X}[_{\mathcal{P}}^{log}$ such that $\{W,]\bar{X}[_{\mathcal{P}}^{log} -]X[_{\mathcal{P}}^{log}\}$ is an admissible covering of $]\bar{X}[_{\mathcal{P}}^{log}$. Given a sheaf of \mathcal{O}_W modules \mathcal{E} we define the sheaf of overconvergent sections as the sheaf $j_W^{\dagger} \mathcal{E} = \varinjlim_{W'} \alpha_{W',]\bar{X}[_{\mathcal{P}}^{log}}^{-1} \alpha_{W', W}^{-1} \mathcal{E}$, where W' varies among the strict neighborhood of $]X[_{\mathcal{P}}^{log}$ in $]\bar{X}[_{\mathcal{P}}^{log}$ that are contained in W and $\alpha_{T, T'} : T \hookrightarrow T'$ is the natural inclusion. If $W =]\bar{X}[_{\mathcal{P}}^{log}$, then we will denote the sheaf of overconvergent sections by $j^{\dagger} \mathcal{E}$.

We suppose that there exists a commutative diagram

$$\begin{array}{ccc} (\bar{X}, M_{\bar{X}}) & \xrightarrow{j} & (\mathcal{P}, M_{\mathcal{P}}) \\ \downarrow g & & \downarrow h \\ (\bar{Y}, M_{\bar{Y}}) & \xrightarrow{i} & (\mathcal{Y}, M_{\mathcal{Y}}) \end{array} \quad (24)$$

where j is a closed immersion and h is formally log smooth. If we denote by $(\mathcal{P}(1), M_{\mathcal{P}(1)})$ (resp. $(\mathcal{P}(2), M_{\mathcal{P}(2)})$) the fiber product of $(\mathcal{P}, M_{\mathcal{P}})$ with itself over $(\mathcal{Y}, M_{\mathcal{Y}})$ (reps. the fiber product of $(\mathcal{P}, M_{\mathcal{P}})$ with itself over $(\mathcal{Y}, M_{\mathcal{Y}})$ three times), then the projections and the diagonal induce the following maps:

$$\begin{aligned} p_i :]\bar{X}[_{\mathcal{P}(1)}^{log} &\rightarrow]\bar{X}[_{\mathcal{P}}^{log} \text{ for } i = 1, 2, \\ p_{i,j} :]\bar{X}[_{\mathcal{P}(2)}^{log} &\rightarrow]\bar{X}[_{\mathcal{P}(1)}^{log} \text{ for } 1 \leq i, j \leq 3, \\ \Delta :]\bar{X}[_{\mathcal{P}}^{log} &\rightarrow]\bar{X}[_{\mathcal{P}(1)}^{log}. \end{aligned}$$

Definition 42. With the previous notation a log overconvergent isocrystal is a pair (\mathcal{E}, ϵ) consisting of a coherent $j^\dagger \mathcal{O}_{\bar{X}[\mathcal{P}]^{\log}}$ -module \mathcal{E} and ϵ a $j^\dagger \mathcal{O}_{\bar{X}[\mathcal{P}(1)]^{\log}}$ -linear isomorphism $\epsilon : p_1^* \mathcal{E} \rightarrow p_2^* \mathcal{E}$ that satisfies $\Delta^*(\epsilon) = \text{id}$ and $p_{1,2}^*(\epsilon) \circ p_{2,3}^*(\epsilon) = p_{1,3}^*(\epsilon)$. We denote by $I^\dagger(((X, \bar{X})/\mathcal{Y}, \mathcal{P})^{\log})$ the category of log overconvergent isocrystals on $((X, M_X), (\bar{X}, M_{\bar{X}})/\mathcal{Y})$ over $(\mathcal{P}, M_{\mathcal{P}})$. We say that (\mathcal{E}, ϵ) is a locally free log overconvergent isocrystal if \mathcal{E} is a locally free $j^\dagger \mathcal{O}_{\bar{X}[\mathcal{P}]^{\log}}$ -module and we indicate the category of locally free log overconvergent isocrystals with $I^\dagger(((X, \bar{X})/\mathcal{Y}, \mathcal{P})^{\log})^{lf}$.

In the case of trivial log structures the previous definition coincides with the definition of overconvergent isocrystals given by Berthelot [Be].

Remark 43. Shiho in [Sh4] definition 4.2 defines the category of log overconvergent isocrystals also in a more general situation, but for our purposes the definition we gave is sufficient.

Given a log pair $((X, M_X), (\bar{X}, M_{\bar{X}}))$, we assume the existence of a diagram

$$(\bar{X}, M_{\bar{X}}) \xrightarrow{g} (\bar{Y}, M_{\bar{Y}}) \xrightarrow{i} (\mathcal{Y}, M_{\mathcal{Y}}), \quad (25)$$

where $(\bar{Y}, M_{\bar{Y}})$ is a log scheme over (S, M_S) , $(\mathcal{Y}, M_{\mathcal{Y}})$ is a p -adic log formal scheme over $(\mathcal{S}, M_{\mathcal{S}})$ and i is a closed immersion.

Coming back to the setting discussed in section 7, we consider the log triple $((U_k, M), (X_k, M), (\hat{X}, M))$ over $((\text{Spec}(k), N), (\text{Spec}(k), N), (\text{Spf}(V), N))$ and the following commutative diagram

$$\begin{array}{ccccc} (U_k, M) & \hookrightarrow & (X_k, M) & \hookrightarrow & (\hat{X}, M) \\ \downarrow & & \downarrow f_k & & \downarrow \\ (\text{Spec}(k), N) & \hookrightarrow & (\text{Spec}(k), N) & \hookrightarrow & (\text{Spf}(V), N) \end{array}$$

The diagram as in (25) is given by

$$(U_k, M) \xrightarrow{g} (\text{Spec}(k), N) \xrightarrow{i} (\text{Spf}(V), N)$$

and the commutative diagram as in (24) is

$$\begin{array}{ccc} (X_k, M) & \xrightarrow{j} & (\hat{X}, M) \\ \downarrow g & & \downarrow h \\ (\text{Spec}(k), N) & \xrightarrow{i} & (\text{Spf}(V), N) \end{array}$$

Let us note that since the immersion $(U_k, M) \hookrightarrow (\hat{X}, M)$ is strict and the closed immersion $(X_k, M) \hookrightarrow (\hat{X}, M)$ is exact, the log tubes in these cases coincide with the classical tubes:

$$]X_k[_{\hat{X}}^{\log} =]X_k[_{\hat{X}} = \hat{X}_K$$

$$]U_k[_{\hat{X}}^{\log} =]U_k[_{\hat{X}} = \hat{U}_K.$$

Now we want to give a description of integrable connections associated to locally free overconvergent isocrystals in our case. By proposition 2.1.10 of [Be] we know that there is an equivalence of categories between $\text{Coh}(j^\dagger \mathcal{O}_{X_k[\hat{X}]})$, the category of $j^\dagger \mathcal{O}_{X_k[\hat{X}]}$ -coherent modules, and the inductive limit category of coherent modules over strict neighborhoods of $]U_k[_{\hat{X}}$ in $]X_k[_{\hat{X}}$. Thanks to remark after proposition 2.1.10 of [Be], if (\mathcal{E}, ϵ) is a locally free log overconvergent isocrystal, then \mathcal{E} is a locally free $j^\dagger \mathcal{O}_{X_k[\hat{X}]}$ -module, which means that there exists a strict neighborhood W of $]U_k[_{\hat{X}}$ in $]X_k[_{\hat{X}}$ and a locally free \mathcal{O}_W -module E , such that

$$j_W^\dagger E = \mathcal{E}.$$

The log overconvergent isocrystal (\mathcal{E}, ϵ) induces an integrable connection on \mathcal{E}

$$\nabla : \mathcal{E} \rightarrow \mathcal{E} \otimes_{j^\dagger \mathcal{O}_{|X_k|_{\hat{X}}}} j^\dagger \omega_{|X_k|_{\hat{X}}/K}^1$$

where $\omega_{|X_k|_{\hat{X}}/K}^1$ is the restriction of $K \otimes \omega_{(\hat{X}, M)/(\mathrm{Spf}(V), N)}^1$ to $|X_k|_{\hat{X}}$; moreover given a strict neighborhood W of $|U_k|_{\hat{X}}$ in $|X_k|_{\hat{X}}$, as we saw before, there exists E on W such that $j_W^\dagger E = \mathcal{E}$ and there exists an integrable connection

$$\nabla : E \rightarrow E \otimes \omega_{W/K}^1$$

which induces the above connection on \mathcal{E} , where $\omega_{W/K}^1$ is the restriction of $K \otimes \omega_{(\hat{X}, M)/(\mathrm{Spf}(V), N)}^1$ to W . If étale locally we are in the situation described in (16), then W contains a subspace of the form

$$\{p \in \hat{X}_K \mid \forall j \ |y_j(p)| \geq \lambda\}$$

for some $\lambda \in (0, 1) \cap \Gamma^*$ with Γ^* the divisible closure of the image of the absolute value $|\cdot| : K^* \rightarrow \mathbb{R}^+$. Therefore we can restrict E to the space

$$\{p \in \hat{X}_{i,j;K}^\circ \mid \lambda \leq |y_j(p)| < 1\}.$$

Proposition 44. *There is an isomorphism*

$$\phi : \hat{D}_{j,i;K}^\circ \times A_K^1([\lambda, 1)) \rightarrow \{p \in \hat{X}_{i,j;K}^\circ \mid \lambda \leq |y_j(p)| < 1\},$$

where $A_K^1([\lambda, 1)) := \{t \in \mathbb{A}_K^1 \mid |t| \in [\lambda, 1)\}$.

Proof. If we can prove that $\hat{D}_{j,i;K}^\circ \times A_K^1([0, 1)) \cong \hat{X}_{i,j;K}^\circ$, then the isomorphism of the proposition will be clear. To prove this we will apply lemma 4.3.1 of [Ke]: we consider $A = \Gamma(\hat{X}_{i,j;K}^\circ, \mathcal{O}_{X_K})$, and as B the ring $A/y_j A = \Gamma(\hat{D}_{i,j;K}^\circ, \mathcal{O}_{X_K})$. We can apply lemma 4.3.1 of [Ke] because $\Gamma(\hat{D}_{i,j;K}^\circ, \mathcal{O}_{X_K})$ is formally smooth over K and we can conclude that

$$\Gamma(\hat{X}_{i,j;K}^\circ, \mathcal{O}_{X_K}) \cong \Gamma(\hat{D}_{i,j;K}^\circ, \mathcal{O}_{X_K})[[y_j]]$$

i.e. the isomorphism that we wanted. \square

We fix a set $\Sigma = \prod_{h=1}^k \Sigma_h \in \mathbb{Z}_p^k$, where k is the number of the irreducible components of $\hat{D} = \bigcup_{h=1}^k \hat{D}^h$ in \hat{X} , with the same notations as before definition 35.

Definition 45. *A log overconvergent isocrystal \mathcal{E} has Σ -unipotent monodromy if there exists an étale covering $\coprod_l \phi_l : \coprod_l \hat{X}_l \rightarrow \hat{X}$ such that every \hat{X}_l has a diagram*

$$\begin{array}{ccc} \hat{D}_l = \bigcup_{j=1}^s \hat{D}_{j,l} & \hookrightarrow & \hat{X}_l \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y_{j,l} = 0\} & \hookrightarrow & \mathrm{Spf} V \{x_{1,l}, \dots, x_{n,l}, y_{1,l}, \dots, y_{m,l}\} / (x_{1,l} \dots x_{r,l} - \pi) \end{array} \quad (26)$$

as in (16) with $\hat{D}_l := \phi_l^{-1}(\hat{D})$ such that for every l the restriction of the log- ∇ -module E_l on $\hat{X}_{l;K}$ to

$$\hat{D}_{i,j,l;K}^\circ \times A_K^1([\lambda, 1)),$$

is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent, $\forall (i, j)$ such that $\phi_l(\hat{D}_{i,j,l}^\circ) \subset \hat{D}^{h(i,j)}$.

We will denote by $I^\dagger((U_k, M), (X_k, M))/(\mathrm{Spf}(V), N)^\Sigma = I^\dagger(((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma})$ the category of log overconvergent isocrystals with Σ -unipotent monodromy.

Remark 46. In definition 45 we do not ask any locally freeness hypothesis, because every object in the category $I^\dagger((U_k, M), (X_k, M))/(\mathrm{Spf}(V), N)$ is such that \mathcal{E} is locally free. This is clear because (\mathcal{E}, ϵ) induces on a strict neighborhood W an \mathcal{O}_W -module E , such that $j_W^\dagger E = \mathcal{E}$ endowed with an integrable connection. As K is of characteristic 0 we can conclude that E is locally free and \mathcal{E} is locally free.

11 Unipotence, generization and overconvergent generization

In this section we recall the three propositions that we will use in the proof of the main theorem. They are proven by Shiho in [Sh6] as generalization of the analogous propositions proven by Kedlaya in [Ke] (assuming $\Sigma = 0$). We will write the statements only in the cases that we need, that are simplified versions of the propositions given in paragraph 2 of [Sh6]. The first property that we consider is called by Shiho and Kedlaya generization property for monodromy ([Ke] proposition 3.4.3 and [Sh6] proposition 2.4).

Proposition 47. *Let A be an affinoid algebra such that $Y = \mathrm{Spm}(A)$ is smooth and endowed with sections y_1, \dots, y_s that are smooth and meet transversally. Suppose that there exists $A \subseteq L$ such that L is an affinoid algebra over K , $\mathrm{Spm}(L)$ is smooth, all the y_i 's are invertible in L and the spectral norm on L restricts to the spectral norm on A . Let I be a quasi open interval contained in $[0, 1)$ and $A_L^n(I)$ defined as $\mathrm{Spm}(L) \times A_K^n(I)$. Let $\Sigma \subset \mathbb{Z}_p^{n+s}$ be a set which is (NLD) and (NID); if $E \in \mathrm{LNM}_{Y \times A_K^n(I), \Sigma}$ is such that the induced object $F \in \mathrm{LNM}_{A_L^n(I), \Sigma}$ is unipotent, then E is Σ -unipotent.*

The second result that we need is called overconvergent generization and describes the property of extension of unipotence on strict neighborhoods (proposition 2.7 of [Sh6] and proposition 3.5.3 of [Ke]).

Proposition 48. *Let P be a p -adic formal affine scheme topologically of finite type over V . Let $Y_k \subseteq P_k$ be an open dense subscheme of the special fiber of P such that P is formally smooth over V in a neighborhood of Y_k . Let W be a strict neighborhood of $]Y_k[_P$ in P_K , $I \subset [0, 1)$ a quasi open interval and Σ a subset of \mathbb{Z}_p^n . Given $E \in \mathrm{LNM}_{W \times A_K^n(I), \Sigma}$ such that the restriction of E to $]Y_k[_P \times A_K^n(I)$ is Σ -unipotent, then for every closed interval $[b, c] \subseteq I$ there exists a strict neighborhood W' of $]Y_k[_P$ in P_K such that W' is contained in W and such that the restriction of E to $W' \times A_K^n[b, c]$ is Σ -unipotent.*

The third property that we need states that, under certain assumptions, a log- ∇ -module with exponents in Σ that is convergent is Σ -unipotent (proposition 2.12 of [Sh6] and lemma 3.6.2 of [Ke]).

Proposition 49. *Let A be an integral affinoid algebra such that $Y = \mathrm{Spm}(A)$ is smooth. Suppose that there exists $A \subseteq L$ such that L is an affinoid algebra over K , $\mathrm{Spm}(L)$ is smooth and y_j 's are invertible in L and the spectral norm of L restricts to the spectral norm of A . Let $\Sigma \subset \mathbb{Z}_p^{n+s}$ be a set which is (NLD) and (NID); if E is an object of $\mathrm{LNM}_{Y \times A_K^n([0,1)), \Sigma}$ which is log convergent ([Sh6], definition 2.9), then it is Σ -unipotent.*

12 Extension theorem

Now we come back to the semistable situation and we will prove that the definition of Σ -unipotent monodromy is well posed.

Proposition 50. *Let \mathcal{E} be an overconvergent log isocrystal which is in the category $I^\dagger((U_k, X_k)/\mathrm{Spf}(V))^{log, \Sigma}$. The notion of Σ -unipotent monodromy for \mathcal{E} is independent of the choice of the étale cover and of the diagram in (16) which we have chosen in definition 45.*

Proof. First we will prove that if \mathcal{E} has Σ -unipotent monodromy for some diagram as in (16), then it has Σ -unipotent monodromy for any diagram as in (16). So we suppose that there exists an étale covering $\coprod_l \phi_l : \coprod_l \hat{X}_l \rightarrow \hat{X}$ such that every \hat{X}_l has a diagram as in (16). As we saw before \mathcal{E} induces on some W strict neighborhood of $]U_k[_{\hat{X}}$ in $]X_k[_{\hat{X}}$ a locally free \mathcal{O}_W -module E with an integrable connection. In the situation of (16) W contains the set

$$\{p \in \hat{X}_{i,K} \mid \forall j \ |y_j(p)| \geq \lambda\}$$

and the restriction of E to

$$\hat{D}_{i,j,K}^\circ \times A_K^1[\lambda, 1)$$

is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent for some λ , if (i, j) are such that $\phi_l(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

Using theorem 40 we can extend E to a module with connection on

$$\hat{D}_{i,j,K}^\circ \times A_K^1[0, 1)$$

that is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent. This is true for every (i, j) so E can be extended to a locally free isocrystal on $((\coprod_{(i,j)} \hat{X}_{i,j}^\circ, M)/(\mathrm{Spf}(V), N))$ which is convergent because it is convergent on $\hat{U} \cap \coprod_{(i,j)} \hat{X}_{i,j}^\circ$ that is an open dense $\hat{X} \cap \coprod_{(i,j)} \hat{X}_{i,j}^\circ$ (we are applying here proposition 28). Hence we have a locally free isocrystal on $((\hat{X}^\circ, M)/(\mathrm{Spf}(V), N))$ which is convergent and such that has exponents along \hat{D}_j in $\cap_{i=1}^r \Sigma_{h(i,j)}$ for every (i, j) such $\phi_l(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

For any other diagram as in (16), E on $\hat{D}_{i,j;K}^\circ \times A_K^1[\lambda, 1]$ with (i, j) such that $\phi_l(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$ is the restriction of a convergent module with connection on $\hat{D}_{i,j;K}^\circ \times A_K^1[0, 1]$ with exponents in $\cap_{i=1}^r \Sigma_{h(i,j)}$, so that it is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on $\hat{D}_{i,j;K}^\circ \times A_K^1[\lambda, 1]$ by proposition 49.

Now we prove that the notion of Σ -unipotence is independent of the choice of the étale covering. To do this, it suffices to prove that if \mathcal{E} is a log overconvergent isocrystal with Σ -unipotent monodromy, for any étale morphism $\phi : \hat{X}' \rightarrow \hat{X}$ such that \hat{X}' admits a diagram

$$\begin{array}{ccc} \hat{D}' = \bigcup_{j=1}^s \hat{D}'_j & \hookrightarrow & \hat{X}' \\ \downarrow & & \downarrow \\ \bigcup_{j=1}^s \{y'_j = 0\} & \hookrightarrow & \mathrm{Spf} V \{x'_1, \dots, x'_n, y'_1, \dots, y'_m\} / (x'_1 \dots x'_r - \pi) \end{array} \quad (27)$$

as in (16), with $\hat{D}' := \phi^{-1}(\hat{D})$, the log- ∇ -module E induced on \hat{X}' by \mathcal{E} is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on $\hat{D}'_{i,j;K}^\circ \times A_K^1[\lambda, 1]$, for every (i, j) such that $\hat{D}'_{i,j}^\circ$ is such that $\phi(\hat{D}'_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$. We may assume that $\hat{D}'_{i,j;K}^\circ = \mathrm{Spm} A$ is affinoid. As in the proof of proposition 36 we know that there exists an étale covering $\coprod_l \hat{X}'_l \rightarrow \hat{X}'$ such that, for any l , \hat{X}'_l admits a diagram as in (16) such that \mathcal{E} has Σ -unipotent monodromy with respect to this diagram.

However since we have already showed that the notion of Σ -unipotent monodromy does not depend on the choice of a diagram as in (16), we can say that \mathcal{E} has Σ -unipotent monodromy with respect to a diagram as in (16) for \hat{X}'_l induced by the diagram (27). This means that the log- ∇ -module E is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent when it is restricted to $\coprod_l (\hat{D}'_{i,j;K}^\circ \times_{\hat{X}_K} \hat{X}'_{l,K}) \times A_K^1[\lambda, 1]$ with (i, j) such that $\phi(\coprod_l (\hat{D}'_{i,j}^\circ \times_{\hat{X}} \hat{X}'_l)) \subset \hat{D}^{h(i,j)}$. Let us take an affine covering $\coprod_h \hat{C}_h \rightarrow \coprod_l (\hat{D}'_{i,j}^\circ \times_{\hat{X}} \hat{X}'_l)$ by affine formal schemes and put $\mathrm{Spm} L := \coprod_h \hat{C}_{h;K}$. Then E is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on $\mathrm{Spm} L \times A_K^1[\lambda, 1]$ and the affinoid algebras A and L satisfy the assumption of proposition 47. Applying proposition 47 we can conclude that E is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on $\hat{D}'_{i,j;K}^\circ \times A_K^1[\lambda, 1]$ as we wanted. \square

We can now state the main result, an extension theorem that generalizes theorem 6.4.5 of [Ke] and theorem 3.16 of [Sh6]. The strategy of the proof is the same as the one in [Ke] and [Sh6] and we will follow step by step the proof of theorem 3.16 of [Sh6].

We will need for the proof the following lemma:

Lemma 51. *If W is an open dense in $P = \mathrm{Spf}(A)$, with A a formal V algebra of topologically finite type such that A_k is reduced, then the spectral seminorm on $\mathcal{O}(\mathbb{I}W_k)_P$ restricts to the spectral seminorm on $\mathcal{O}(\mathbb{I}P_k)_P$.*

Proof. We can suppose that W is defined by the equation $\{g \neq 0\}$, in particular that $W = \mathrm{Spf}(A \left\{ \frac{1}{g} \right\})$.

So we have a map of V -algebras

$$A \longrightarrow A \left\{ \frac{1}{g} \right\},$$

which is injective modulo π because W_k is dense in A_k that is reduced. By the topological Nakayama's lemma (ex 7.2 of [Ei]) we can conclude that we have an inclusion of V -algebras

$$A \hookrightarrow A \left\{ \frac{1}{g} \right\}$$

which induces an inclusion of affinoid algebras

$$\mathcal{O}(]P_k[_P) = A \otimes K \hookrightarrow \mathcal{O}(]W_k[_P) = A \left\{ \frac{1}{g} \right\} \otimes K$$

Let us take the Banach norm $| \cdot |_P$ on $A \otimes K$ induced by A and the Banach norm $| \cdot |_W$ on $A \left\{ \frac{1}{g} \right\} \otimes K$ induced by $A \left\{ \frac{1}{g} \right\}$, then

$$|a|_P = |a|_W$$

for any $a \in K \otimes A$. By the well known formula (see for example [FvdP] corollary 3.4.6)

$$|a|_{P,\text{sp}} = \lim_{n \rightarrow \infty} |a^n|_P^{\frac{1}{n}}$$

where with $| \cdot |_{P,\text{sp}}$ we denote the spectral norm, we are done. \square

Theorem 52. *We fix a set $\Sigma = \prod_{h=1}^k \Sigma_h \in \mathbb{Z}_p^k$, where k is the number of the irreducible components of $\hat{D} = \bigcup_{h=1}^k \hat{D}^h$ in \hat{X} and we require that Σ has the properties (NID) and (NLD). Let us suppose that locally for the étale topology we have a diagram as (16), then the restriction functor*

$$j^\dagger : I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))^\Sigma \longrightarrow I^\dagger((U_k, M), (X_k, M))/(\text{Spf}(V), N)^\Sigma$$

is an equivalence of categories.

Proof. We will divide the proof in 3 steps.

Step 1: the functor j^\dagger is well defined.

Let \mathcal{E} be in the category $I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))^\Sigma$, then we have to prove that $j^\dagger(\mathcal{E})$ is in the category $I^\dagger((U_k, M), (X_k, M))/(\text{Spf}(V), N)^\Sigma$. Thanks to lemma 36 and lemma 50, we can work étale locally. We suppose that ϕ is an étale map to \hat{X} and we call again \hat{X} an étale neighborhood for which we have the diagram as in (16); in this situation the log convergent isocrystal \mathcal{E} induces a log- ∇ -module with respect to y_1, \dots, y_s on \hat{X}_K such that has exponents along $\hat{D}_{j;K}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$ if i and j are such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$. By the definition of Σ -unipotent monodromy (definition 45) we are reduced to prove that, if we restrict E to

$$\hat{D}_{i,j;K}^\circ \times A_K^1[0, 1),$$

then it is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent if i, j are such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

We know by hypothesis that the restriction of E to $\hat{D}_{i,j;K}^\circ \times A_K^1[0, 1)$ is log convergent and has exponents along $\hat{D}_{j;K}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$, hence we have to prove that this implies $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotence; from proposition 49 we know that this implies $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotence.

Step 2: the functor j^\dagger is fully faithful.

We have to prove that given $f : \mathcal{E} \rightarrow \mathcal{F}$, a morphism of log overconvergent isocrystals of Σ -unipotent monodromy, if there exist extensions of \mathcal{E} and \mathcal{F} to log convergent isocrystals with exponents in Σ that we call respectively $\tilde{\mathcal{E}}$ and $\tilde{\mathcal{F}}$, then f extends uniquely to $\tilde{f} : \tilde{\mathcal{E}} \rightarrow \tilde{\mathcal{F}}$.

We can work étale locally. We denote by ϕ an étale map to \hat{X} and again by \hat{X} an étale neighborhood that we consider for which there exists a diagram as in (16).

Let us take W a strict neighborhood of $]U_k[_{\hat{X}}$ in $]X_k[_{\hat{X}}$; by definition f induces a morphism φ of ∇ -modules between $E_{\mathcal{E}}$ and $E_{\mathcal{F}}$, the ∇ -modules on W that are induced by \mathcal{E} and \mathcal{F} respectively:

$$\varphi : E_{\mathcal{E}} \rightarrow E_{\mathcal{F}}.$$

We call $E_{\tilde{\mathcal{E}}}$ and $E_{\tilde{\mathcal{F}}}$ the log- ∇ -modules on \hat{X}_K induced by $\tilde{\mathcal{E}}$ and $\tilde{\mathcal{F}}$.

Let us take the following covering of

$$\hat{X}_K = \bigcup_{J \subset \{1, \dots, s\}} A_J$$

where

$$A_J = \{p \in \hat{X}_K \mid |y_j(p)| < 1 \ (j \in J) \ |y_j(p)| \geq \lambda \ (j \notin J)\}$$

and $\lambda \in (0, 1) \cup \Gamma^*$ is such that both \mathcal{E} and \mathcal{F} are defined on the following set:

$$B = \{p \in \hat{X}_K \mid |y_j(p)| \geq \lambda \ \forall j\}.$$

The covering of \hat{X}_K given by the A_J 's restricts to the following covering of $B = \bigcup_{J \subset \{1, \dots, s\}} B_J$, where

$$B_J = \{p \in \hat{X}_K \mid \lambda \leq |y_j(p)| < 1 \ (j \in J), \ |y_j(p)| \geq \lambda \ (j \notin J)\}.$$

The extensions $E_{\mathcal{E}}$ and $E_{\mathcal{F}}$ are log convergent in

$$\{p \in \hat{X}_K \mid y_j(p) = 0 \ (j \in J), \ |y_j(p)| \geq \lambda \ (j \notin J)\} \times A^{|J|}[0, 1) \quad (28)$$

by proposition 3.6 of [Sh6] and they have exponents in $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$. They extend the restrictions of $E_{\mathcal{E}}$ and $E_{\mathcal{F}}$ on

$$\{p \in \hat{X}_K \mid y_j(p) = 0 \ (j \in J) \ |y_j(p)| \geq \lambda \ (j \notin J)\} \times A^{|J|}[\lambda, 1).$$

By theorem 40 we can conclude that ϕ extends to

$$A_J = \{p \in \hat{X}_K \mid y_j(p) = 0 \ (j \in J) \ |y_j(p)| \geq \lambda \ (j \notin J)\} \times A^{|J|}[0, 1);$$

this means that on this set there exists a unique

$$\tilde{\phi}_J : E_{\mathcal{E}} \rightarrow E_{\mathcal{F}}$$

that extends ϕ on B_J .

On $A_I \cap A_J$ we have the extensions ϕ_I and ϕ_J , which glue because they coincide on the set

$$B_I \cap B_J = \{p \in \hat{X}_K \mid \lambda \leq |y_j(p)| < 1 \ (j \in (I \cup J) - (I \cap J)), \\ |y_j(p)| \geq \lambda \ (j \notin (I \cup J))\} \times A^{|I \cap J|}[\lambda, 1)$$

because they extend the map ϕ on $B_I \cap B_J$.

Step 3: the functor j^\dagger is essentially surjective.

Since we have the étale descent property for the category of locally free log convergent isocrystals (remark 5.1.7 of [Sh1]) and the full faithfulness of the functor j^\dagger , we may work étale locally to prove the essential surjectivity.

If $\mathcal{E} \in I^\dagger((U_k, X_k), \mathrm{Spf}(V))^{log, \Sigma}$, then by definition of log overconvergent isocrystal we know that \mathcal{E} induces a module with connection on the following set

$$\{p \in \hat{X}_K \mid \forall j : |y_j(p)| \geq \lambda\}$$

that we will denote by E that is $\cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on

$$\hat{D}_{i,j,K}^\circ \times A^1[\lambda, 1) = \{p \in \hat{D}_{j,K} \mid \forall j' \neq j : |y_{j'}(p)| = 1, \ \forall i' \neq i : |x_{i'}(p)| = 1\} \times A^1[\lambda, 1),$$

for i, j such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

We will prove that E extends to a log- ∇ -module on $C_{a,\lambda} = \{p \in \hat{X}_K \mid \forall j > a, |y_j(p)| \geq \lambda\} \ \forall a = 0, \dots, s$ such that has exponents along $\{y_j = 0\}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$ with i, j such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$, proceeding by induction on a .

So we suppose, by induction hypothesis, that E extends to the set $C_{a-1,\lambda} = \{p \in \hat{X}_K \mid \forall j > a-1, |y_j(p)| \geq \lambda\}$ for some λ with exponents along $\{y_j = 0\}$ in $\cap_{i=1}^r \Sigma_{h(i,j)}$ with i, j such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^{h(i,j)}$.

We consider the following admissible covering of $\hat{X}_K = A \cup B$, where

$$A = \{p \in \hat{X}_K \mid |y_a(p)| \geq \lambda'\}$$

$$B = \{p \in \hat{X}_K \mid |y_a(p)| < 1\}$$

with $\lambda' \in [\lambda, 1) \cap \Gamma^*$.

Intersecting the covering $A \cup B$ with $C_{a-1, \lambda'}$ we obtain the following admissible covering:

$$\begin{aligned} C_{a-1, \lambda'} &= (C_{a-1, \lambda'} \cap A) \cup (C_{a-1, \lambda'} \cap B) = \\ &= \{p \in \hat{X}_K \mid \forall j > a-1 \mid |y_j(p)| \geq \lambda'\} \cup \\ &\quad \{p \in \hat{X}_K \mid \forall j > a \mid |y_j(p)| \geq \lambda', \lambda' \leq |y_a(p)| < 1\} = \\ &= \{p \in \hat{X}_K \mid \forall j > a-1 \mid |y_j(p)| \geq \lambda'\} \cup \\ &\quad \{p \in \hat{D}_{a, K} \mid \forall j > a \mid |y_j(p)| \geq \lambda'\} \times A^1[\lambda', 1), \end{aligned} \tag{29}$$

and intersecting with $C_{a, \lambda'}$:

$$\begin{aligned} C_{a, \lambda'} &= (C_{a, \lambda'} \cap A) \cup (C_{a, \lambda'} \cap B) = \\ &= \{p \in \hat{X}_K \mid \forall j > a-1 \mid |y_j(p)| \geq \lambda'\} \cup \{p \in \hat{X}_K \mid \forall j > a \mid |y_j(p)| \geq \lambda', |y_a(p)| < 1\} = \\ &= \{p \in \hat{X}_K \mid \forall j > a-1 \mid |y_j(p)| \geq \lambda'\} \cup \{p \in \hat{D}_{a, K} \mid \forall j > a \mid |y_j(p)| \geq \lambda'\} \times A^1[0, 1). \end{aligned} \tag{30}$$

Comparing the formulas in (29) and (30), we see that it is sufficient to prove that E extends from

$$\{p \in \hat{D}_{a, K} \mid \forall j > a, |y_j(p)| \geq \lambda'\} \times A_K^1[\lambda', 1)$$

to

$$\{p \in \hat{D}_{a, K} \mid \forall j > a \mid |y_j(p)| \geq \lambda'\} \times A^1[0, 1),$$

for some $\lambda' \in [\lambda, 1) \cap \Gamma^*$ in a log- ∇ -module such that it has exponents along $\{y_j = 0\}$ in $\cap_{i=1}^r \Sigma_{h(i, j)}$ with i, j for which $\phi(\hat{D}_{i, j}^\circ) \subset \hat{D}^{h(i, j)}$.

As we saw before E is $\cap_{i=1}^r \Sigma_{h(i, a)}$ -unipotent on

$$\begin{aligned} \hat{D}_{i, a, K}^\circ \times A_K^1([\lambda, 1)) &= \\ \{p \in \hat{D}_{a, K} \mid \forall j' \neq a : |y_{j'}'(p)| = 1, \forall i' \neq i : |x_{i'}'(p)| = 1\} \times A_K^1([\lambda, 1)), \end{aligned} \tag{31}$$

so it is $\cap_{i=1}^r \Sigma_{h(i, a)}$ -unipotent also on

$$\begin{aligned} \prod_i \hat{D}_{i, a, K}^\circ \times A_K^1([\lambda, 1)) &= \\ \prod_i \{p \in \hat{D}_{a, K} \mid \forall j' \neq a : |y_{j'}'(p)| = 1, \forall i' \neq i : |x_{i'}'(p)| = 1\} \times A_K^1([\lambda, 1)). \end{aligned}$$

We now want to apply proposition 48; following the notation given in the proposition in our case we have that P is the pull-back of

$$\mathrm{Spf} \frac{V\{x_1, \dots, x_n, y_1 \dots \hat{y}_a \dots y_m\}}{x_1 \dots x_r - \pi} \{y_j^{-1} \mid j < a\} \left\{ \prod_{i, i' \atop i \neq i'} (x_i - x_{i'})^{-1} \right\}$$

by the morphism

$$\hat{X} \rightarrow \mathrm{Spf} V\{x_1, \dots, x_n, y_1, \dots, y_m\} / (x_1 \dots x_r - \pi),$$

Y_k is the open defined in P_k by the following equation

$$\{y_{a+1} \dots y_s \neq 0\}$$

and

$$\begin{aligned} W \times A_K^1(I) &= \{p \in \hat{D}_{a, K} \mid \forall j < a : |y_j(p)| = 1, \\ &\quad \forall j > a, |y_j(p)| \geq \lambda \mid \forall i' \neq i : |x_{i'}'(p)| = 1\} \times A_K^1([\lambda, 1)). \end{aligned}$$

The hypothesis of proposition 48 are fulfilled.

The restriction of E to

$$\{p \in \hat{D}_{a,K} \mid \forall j > a, |y_j(p)| \geq \lambda\} \times A_K^1([\lambda, 1))$$

is a log- ∇ -module with exponents in $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$, that is $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on

$$]Y_k[_P = \prod_i \{p \in \hat{D}_{a,K} \mid \forall j' \neq a : |y_{j'}(p)| = 1, \forall i' \neq i : |x_{i'}(p)| = 1\} \times A_K^1([\lambda, 1));$$

so applying proposition 48 we know that for every $[b, c] \subset [\lambda, 1)$ there exists a λ' (we suppose that it verifies $\lambda' \in (c, 1)$ for gluing reasons) such that E is $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on

$$\prod_i \{p \in \hat{D}_{a,K} \mid \forall j < a : |y_j(p)| = 1, \\ \forall j > a : |y_j(p)| \geq \lambda', \forall i' \neq i : |x_{i'}(p)| = 1\} \times A^1([b, c]).$$

Now we apply proposition 47 with

$$\text{Spm}(L) = \prod_i \{p \in \hat{D}_{a,K} \mid \forall j < a : |y_j(p)| = 1, \\ \forall j > a : |y_j(p)| \geq \lambda', \forall i' \neq i : |x_{i'}(p)| = 1\}$$

and

$$Y = \{p \in \hat{D}_{a,K} \mid \forall j > a : |y_j(p)| \geq \lambda'\};$$

we are in the hypothesis of that proposition thanks to lemma 51, hence we deduce that E is $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent on

$$\{p \in \hat{D}_{a,K} \mid \forall j > a : |y_j(p)| \geq \lambda'\} \times A^1((b, c)).$$

By theorem 40 we see that E can be extended to a $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ -unipotent log- ∇ -module, in particular to a log- ∇ -module with exponents in $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ on

$$\{p \in \hat{D}_{a,K} \mid \forall j > a : |y_j(p)| \geq \lambda'\} \times A^1([0, c)).$$

Now we glue this with E and we obtain a log- ∇ -module with exponents in the set $\prod_j \cap_{i=1}^r \Sigma_{h(i,j)}$ on

$$\{p \in \hat{D}_{a,K} \mid \forall j > a, |y_j(p)| \geq \lambda'\} \times A_K^1([0, 1))$$

as we wanted.

Therefore we have a log- ∇ -module defined on the space \hat{X}_K and we now prove that it is convergent.

We know that the restriction of E to \hat{U}_K is log convergent because it is an extension of an overconvergent log isocrystal on \hat{U}_K , hence it belongs to the category

$$I_{\text{conv}}((\hat{U}, M), (\text{Spf}(V), N))^{lf}.$$

Since \hat{U} is an open dense in \hat{X} , we have a module with log connection defined in the whole space that is convergent on an open dense of the space; we can apply proposition 28 and conclude that E is convergent. \square

13 Main theorem

As we saw in proposition 52 there is an equivalence of categories

$$j^\dagger : I_{\text{conv}}((\hat{X}, M)/(\text{Spf}(V), N))^{lf, \Sigma} \longrightarrow I^\dagger((U_k, X_k)/\text{Spf}(V))^{log, \Sigma}.$$

Now we want to compare this to the category $MIC((X_K, M)/K)^{lf}$ that we defined in definition 3. We can define the notion of exponents also in the algebraic case, giving the analogous definition that we gave before definition 35, replacing the rigid analytic space \hat{X}_K with the algebraic space X_K , the divisor \hat{D}_K with the divisor D_K and the $\mathcal{O}_{\hat{X}_K}$ -module $\omega_{(\hat{X}_K, M)/K}^1$ with the \mathcal{O}_{X_K} -module $\omega_{(X_K, M)/K}^1$.

We fix a set $\Theta = \prod_{p=1}^f \Theta_p \subset \bar{K}^f$, where f is the number of irreducible components of the divisor $D_K = \cup_{p=1}^f D^{p;K}$. We say that (E, ∇) in $MIC((X_K, M_D)/K)$ has residue along D_K in Θ if étale locally there exists a diagram analogous to (16) such that for every l the log- ∇ -module $X_{l;K}$ induced by (E, ∇) has exponents along $D_{j,l;K}$ in $\Theta_{p(j)}$ for every j such that $\phi_{l;K}(\hat{D}_{j,l}) \subset D^{p(j)}$.

We will denote the category of locally free module with integrable log connection with exponents in Θ by $MIC((X_K, M_D)/K)^\Theta$. We can prove as in lemma 36 that the notion of exponents in Θ can be given étale locally and that is independent from the choice of a diagram as in (16). If (E, ∇) is in $MIC((X_K, M_D)/K)$ we restrict locally étale in a situation for which there exists a diagram analogous to (16) for the algebraic setting and we look at the exponents of (E, ∇) along $D_{j;K}$. In particular we consider the log- ∇ -module $(\hat{E}, \hat{\nabla})$ on \hat{X}_K induced by the log infinitesimal locally free isocrystal $\Psi(E, \nabla)$ (where Ψ is the functor defined in proposition 16) and the residue of it along $\hat{D}_{j;K}$. We have a map

$$\text{End}_{\mathcal{O}_{D_{j;K}}}(E|_{D_{j;K}}) \rightarrow \text{End}_{\mathcal{O}_{\hat{D}_{j;K}}}(\hat{E}|_{\hat{D}_{j;K}}),$$

that sends the residue of (E, ∇) along $D_{j;K}$ to the residue of $(\hat{E}, \hat{\nabla})$ along $\hat{D}_{j;K}$. Moreover the map is injective because the map

$$\Gamma(D_{j;K}, \mathcal{O}_{D_{j;K}}) \rightarrow \Gamma(\hat{D}_{j;K}, \mathcal{O}_{\hat{D}_{j;K}})$$

is injective. This means that étale locally (E, ∇) and the log- ∇ -module $(\hat{E}, \hat{\nabla})$ have the same exponents along $D_{j;K}$ and $\hat{D}_{j;K}$ respectively.

The relation between Θ and Σ is as follows.

Given a log infinitesimal isocrystal \mathcal{E} with exponents in Σ then the functor Ψ^{-1} associates to it a module with integrable log connection (E, ∇) such that it has exponents in $\Theta = \prod_{p=1}^f \Theta_p$, where the p -th component Θ_p is given by $\cap_{i=1}^r \Sigma^{h(i,j)}$ where j is such that $\phi_{l,K}(D_{j;K}) \subset D^{p;K}$. Viceversa given a module with integrable log connection (E, ∇) such that it has exponents in Θ , the functor Ψ associates to it a log infinitesimal isocrystal \mathcal{E} with exponents in $\Sigma = \prod_{h=1}^k \Sigma_h$ where the h -th component is given by $\Theta^{p(j)}$ where j is such that $\phi(\hat{D}_{i,j}^\circ) \subset \hat{D}^h$.

From this it follows that the functor Ψ induces an equivalence of categories

$$MIC((X_K, M_D)/(K, \text{triv}))^{lf, \Theta} \longrightarrow I_{inf}((\hat{X}, M)/(\text{Spf} V, N))^{lf, \Sigma}.$$

If we start from a log overconvergent isocrystal \mathcal{E} with Σ -unipotent monodromy as in 45, we apply the equivalence of category given by the functor j^\dagger of theorem 52 and the observations written above, we can conclude that there is fully faithful functor

$$I^\dagger((U_k, M), (X_k, M))/(\text{Spf}(V), N)^\Sigma \longrightarrow MIC((X_K, M_D)/(K, \text{triv}))^{lf, \Theta}.$$

The logarithmic extension theorem of Andr e and Baldassarri (theorem 4.9 of [AnBa]) gives an equivalence of category between $MIC((X_K, M_D)/(K, \text{triv}))^{lf, \Theta}$ and the category of coherent modules with connection on U_K regular along D_K , that we denote by $MIC(U_K/K)^{reg}$ gives us the general result.

Theorem 53. *There is a fully faithful functor*

$$I^\dagger((U_k, M), (X_k, M))/(\text{Spf}(V), N)^\Sigma \longrightarrow MIC(U_K/K)^{reg}.$$

Acknowledgements

This work is a generalization of the author's PhD thesis. It is a pleasure to thank her advisors: prof. Francesco Baldassarri and prof. Atsushi Shiho. Much of this work was done while the author was visiting the University of Tokyo; she wants to thank all the people there for their hospitality. In particular she wants to acknowledge prof. Atsushi Shiho for the helpfulness in guiding her work and for the careful reading of her thesis and prof. Takeshi Saito for many useful discussions. The author would like to express her gratitude also to prof. Bruno Chiarellotto and prof. Marco Garuti for valuable advises.

References

- [AnBa] Y. André, F. Baldassarri, *De Rham Cohomology of Differential Modules on Algebraic Varieties*, Progress in Mathematics **189**, Birkhäuser, 2001. Second edition in preparation.
- [BaCh] F. Baldassarri, B. Chiarellotto, *Formal and p -adic theory of differential systems with logarithmic singularities depending upon parameters*, Duke Math. J., **72**, 241-300, (1993).
- [Be] P. Berthelot, *Cohomologie rigide et cohomologie rigide à support propre. Première partie*, prépublication de l'IRMAR, 96-03, (1993).
- [BeOg] P. Berthelot, A. Ogus, *Notes on crystalline cohomology*, Mathematical Notes, Princeton University Press, (1978).
- [Bo] S. Bosch, *Lectures on Formal and Rigid Geometry*, Preprintreihe des Mathematischen Instituts, Heft 378, Westfälische Wilhelms-Universität, Münster, (2005).
- [ChFo] B. Chiarellotto, M. Fornasiero, *Logarithmic de Rham, Infinitesimal and Betti cohomologies*, J. Math. Sci. Univ. Tokyo, **13**, 205-257, (2006).
- [ChTs] B. Chiarellotto, N. Tsuzuki, *Cohomological descent of rigid cohomology for étale coverings*, Rend. Sem. Mat. Univ. Padova, **109**, 63-215, (2003).
- [CtMeII] G. Christol, Z. Mebkhout, *Sur le théorème de l'indice des équations différentielles p -adiques II*, Annals of math., 2nd Ser., **146**, No. 2, 345-410, (1997).
- [CtMeIV] G. Christol, Z. Mebkhout, *Sur le théorème de l'indice des équations différentielles p -adiques IV*, Invent. math., **143**, 629-672, (2001).
- [De] P. Deligne, *Catégories Tannakiennes*, in Grothendieck Festschrift II, Progress in Mathematics, Birkhauser, pp. 111-195, (1990).
- [DeMi] P. Deligne; J. S. Milne, *Tannakian Categories*, in Hodge cycles, motives, and Shimura varieties, Lecture Notes in Mathematics, 900. Springer-Verlag, Berlin-New York, (1982).
- [DGS] B. Dwork, G. Gerotto, and F. Sullivan, *An Introduction to G -Functions*, Annals of Math. Studies **133**, Princeton University Press, Princeton, (1994).
- [Ei] D.Eisenbud, *Commutative algebra with a view toward algebraic geometry*, **GTM 150**, Springer-Verlag, New York, (1995).
- [FvdP] J. Fresnel, M. van der Put, *Rigid analytic geometry and its applications*, Progress in Mathematics **218**, Birkhäuser, (2004).
- [Gr] A. Grothendieck, *Crystals and the de Rham cohomology of schemes*, in *Dix Exposés sur la Cohomologie des Schemas*, pp. 306-358, North Holland publishing company, (1968).
- [Ha] R.Hartshorne, *Algebraic Geometry*, **GTM 52**, Springer-Verlag, New York, (1977).

- [Ka] K. Kato, *Logarithmic structure of Fontaine-Illusie*, in Algebraic Analysis, Geometry and Number Theory, The John Hopkins Univ. Press, 191-224, (1988).
- [Kz] N. M. Katz, *Nilpotent connection and the monodromy theorem: an application of a result of Turrittin*, Pub. Math. IHES, **39**, 175-232, (1970).
- [Ke] K. S. Kedlaya, *Semistable reduction for overconvergent F -isocrystals, I: Unipotence and logarithmic extensions*, Compositio Math., **143**, 1164-1212, (2007).
- [LS] B. Le Stum, *Rigid Cohomology*, Cambridge Tracts in Mathematics 172, Cambridge University Press, (2007).
- [NaSh] Y. Nakkajima, A. Shiho, *Weight filtration on log crystalline cohomologies of families of open smooth varieties*, Lecture Notes in Mathematics, **1959**, Springer-Verlag, (2008).
- [Og] A. Ogus, *F -isocrystals and de Rham cohomology II - convergent isocrystals*, Duke Math. **51**, 765-850, (1984).
- [Sh1] A. Shiho, *Crystalline fundamental groups I- Isocrystals on log crystalline site and log convergent site*, Journal of Mathematical Sciences, University of Tokyo, **7**, 509-656, (2000).
- [Sh2] A. Shiho, *Crystalline fundamental groups I- Log convergent cohomology and rigid cohomology*, Journal of Mathematical Sciences, University of Tokyo, **9**, 1-163, (2002).
- [Sh3] A. Shiho, *Relative log convergent cohomology and relative rigid cohomology I*, arXiv:0707.1742v3, (2007).
- [Sh4] A. Shiho, *Relative log convergent cohomology and relative rigid cohomology II*, arXiv:0707.1743v2, (2007).
- [Sh5] A. Shiho, *Relative log convergent cohomology and relative rigid cohomology III*, arXiv:0805.3229v1, (2008).
- [Sh6] A. Shiho, *On logarithmic extension of overconvergent isocrystals*, Math. Ann., published online, (2010).
- [Sh7] A. Shiho, *Cut-by-curves criterion for the log extendability of overconvergent isocrystals*, Math. Zeit., published online, (2010).